



# Disconnected cuts in claw-free graphs <sup>☆</sup>

Barnaby Martin <sup>a</sup>, Daniël Paulusma <sup>a,\*</sup>, Erik Jan van Leeuwen <sup>b</sup>

<sup>a</sup> Durham University, Durham, United Kingdom

<sup>b</sup> Utrecht University, Utrecht, the Netherlands

## ARTICLE INFO

### Article history:

Received 27 June 2018

Received in revised form 1 April 2020

Accepted 2 April 2020

Available online 11 May 2020

### Keywords:

Vertex cut

Claw-free graph

Line graph

Circular-arc graph

Graph decomposition

Polynomial-time algorithm

## ABSTRACT

A disconnected cut of a connected graph is a vertex cut that itself also induces a disconnected subgraph. The corresponding decision problem is called DISCONNECTED CUT. This problem is known to be NP-hard on general graphs. We prove that it is polynomial-time solvable on claw-free graphs, answering a question of Ito et al. (TCS 2011). The basis for our result is a decomposition theorem for claw-free graphs of diameter 2, which we believe is of independent interest and builds on the research line initiated by Chudnovsky and Seymour (JCTB 2007–2012) and Hermelin et al. (ICALP 2011). On our way to exploit this decomposition theorem, we characterize how disconnected cuts interact with certain cobipartite subgraphs, and prove two further algorithmic results, namely that DISCONNECTED CUT is polynomial-time solvable on circular-arc graphs and line graphs.

© 2020 Elsevier Inc. All rights reserved.

## 1. Introduction

Graph connectivity is a crucial graph property studied in the context of network robustness. Well-studied notions of connectivity consider for example hamiltonicity, edge-disjoint spanning trees, edge cuts, vertex cuts, etc. In this paper, we study the notion of a *disconnected cut*, which is a vertex set  $U$  of a connected graph  $G$  such that  $G - U$  is disconnected and the subgraph  $G[U]$  induced by  $U$  is disconnected as well. Alternatively, we may say that  $V(G)$  can be partitioned into nonempty sets  $V_1, V_2, V_3, V_4$  such that no vertex of  $V_1$  is adjacent to a vertex of  $V_3$  (that is,  $V_1$  is anti-complete to  $V_3$ ) and  $V_2$  is anti-complete to  $V_4$ ; then both  $V_1 \cup V_3$  and  $V_2 \cup V_4$  form a disconnected cut. See Fig. 1 for an example. The DISCONNECTED CUT problem asks whether a given connected graph  $G$  has a disconnected cut.

The DISCONNECTED CUT problem is intimately connected to at least five other problems studied in the literature. We give a brief overview here, and refer to the related work section for more details. The name DISCONNECTED CUT originates from Fleischner et al. [16], who determined the complexity of partitioning the vertices of a graph into exactly  $k$  *bicliques* (complete bipartite graphs with at least one edge), except for the case  $k = 2$ . For  $k = 2$ , this problem is polynomially equivalent to DISCONNECTED CUT (by taking the complement of the input graph). The DISCONNECTED CUT problem can also be seen as an  $H$ -PARTITION problem for appropriately defined 4-vertex graphs  $H$ . Dantas et al. [9] proved that  $H$ -PARTITION is polynomial-time solvable for each 4-vertex graph  $H$  except for the two cases equivalent to DISCONNECTED CUT. If the input graph has diameter 2, then DISCONNECTED CUT is equivalent to  $C_4$ -COMPACTION [16], which asks for a homomorphism  $f$  from a graph  $G$  to the graph  $C_4$  (the 4-vertex-cycle with a self-loop in each vertex) such that for every  $xy \in E(C_4)$  with

<sup>☆</sup> This paper received support from the Leverhulme Trust (RPG-2016-258). An extended abstract of it appeared in the proceedings of ESA 2018 [29].

\* Corresponding author.

E-mail addresses: barnaby.d.martin@durham.ac.uk (B. Martin), daniel.paulusma@durham.ac.uk (D. Paulusma), e.j.vanleeuwen@uu.nl (E.J. van Leeuwen).




Fig. 1. Graph with disconnected cuts  $V_1 \cup V_3$  and  $V_2 \cup V_4$  (figure originally appeared in [23]).

$x \neq y$  there is an edge  $uv \in E(G)$  with  $f(u) = x$  and  $f(v) = y$ . The diameter-2 case is also equivalent to testing if a graph can be modified to a biclique by a series of edge contractions [23]. The restriction to graphs of diameter 2 is natural, as every graph of diameter 1 has no disconnected and every graph of diameter at least 3 has a disconnected cut [16]. Finally, DISCONNECTED CUT fits in the broad study of vertex cut problems with extra properties on the cut set, such as  $(k, \ell)$ -cuts,  $k$ -cuts, (strict)  $k$ -clique cuts, stable cuts, matching cuts, etc.; see e.g. [24] for an overview.

The above demonstrates that DISCONNECTED CUT is of central importance to understanding many different types of problems, ranging from cut problems to homomorphism and graph contractibility problems. Therefore, there has been broad interest to determine its computational complexity. Indeed, numerous papers [7,9–11,16,22,23,32] asked about its complexity on general graphs. NP-completeness was proven independently in [28] and by Vikas, as announced in [34]. The strong interest in DISCONNECTED CUT also led to a study on graph classes. We know polynomial-time algorithms for many classes, including graphs of bounded maximum degree, graphs not locally connected, graphs with a dominating edge (which include cobipartite graphs and  $P_4$ -free graphs) [16];  $2P_2$ -free graphs, co-spiders, co- $P_4$ -sparse graphs, co-circular arc graphs [7]; apex-minor-free graphs (which generalize planar graphs), chordal graphs [23];  $4P_1$ -free graphs (graphs with independence number at most 4), graphs of bounded treewidth,  $(2P_1 + P_2)$ -free graphs (co-diamond-free graphs),  $(C_5, \overline{P}_5)$ -free graphs, co-planar graphs, co- $(q, q - 4)$ -graphs (for every fixed integer  $q$ ), and  $(C_3 + P_1)$ -free graphs (which contains the class of triangle-free graphs) [10]. The latter is the complement of the well-known class of claw-free graphs (graphs with no induced claw  $K_{1,3}$ ). However, even for several simple graphs  $H$ , and thus comparatively well-structured classes of  $H$ -free graphs, the complexity landscape of DISCONNECTED CUT still contains gaps. Indeed, even for four-vertex graphs  $H$ , we show that precisely one open case remains, namely the case  $H = K_4$ . To prove this result, we need to deal with in particular one non-trivial case, namely, when  $H = K_{1,3}$  (the claw).

Our interest in DISCONNECTED CUT on claw-free graphs is heightened by the close relation of this problem to  $C_r$ -CONTRACTIBILITY. The latter problem is to decide if a graph  $G$  contains the  $r$ -vertex cycle  $C_r$  as a contraction. This problem is NP-complete if  $r \geq 4$  [4] and stays NP-complete for claw-free graphs as long as  $r \geq 6$  [15]. The case  $r \leq 3$  is polynomial-time solvable even for general graphs [4]. Hence, for claw-free graphs this leaves open the cases where  $r \in \{4, 5\}$ . Ito et al. [23] showed that  $C_4$ -CONTRACTIBILITY on claw-free graphs of diameter 2 is equivalent to DISCONNECTED CUT. As DISCONNECTED CUT is trivial if the input graph does not have diameter 2, this led Ito et al. [23] to explicitly ask the following:

*What is the computational complexity of DISCONNECTED CUT on claw-free graphs?*

### 1.1. Our contribution

We answer the open question of Ito et al. [23] by giving a polynomial-time algorithm for DISCONNECTED CUT on claw-free graphs (which also finds a disconnected cut if it exists). This immediately implies that  $C_4$ -COMPACTION and  $C_4$ -CONTRACTIBILITY are polynomial-solvable on claw-free graphs of diameter 2. As claw-free graphs are not closed under edge contraction, the latter is certainly not expected beforehand.

Our result is grounded in a new graph-theoretic theorem that proves that claw-free graphs of diameter 2 belong to one of four basic graph classes after performing two types of elementary operations. The theorem builds on one of the algorithmic decomposition theorems for claw-free graphs developed by Hermelin et al. [21], and relies on the pioneering works of Chudnovsky and Seymour [6]. Several other algorithmic decomposition theorems for claw-free graphs have been built on the ideas of Chudnovsky and Seymour, see e.g. [12,25], which jointly have had a broad impact on our algorithmic understanding of claw-free graphs (see [21] for an overview or [3]). Our structural theorem and resulting algorithm for DISCONNECTED CUT expand this line of research.

The crux of the proof of our structural theorem is to exploit the extra structure offered by claw-free graphs of diameter 2 to show that the so-called strip-structures, which are central to the aforementioned decomposition theorems, only contain trivial strips. An important ingredient in the proof is to exclude not only twins (vertices  $u, v$  for which  $N[u] = N[v]$ ), but also vertices with nested neighbourhoods (vertices  $u$  for which there exists a vertex  $v$  such that  $N(u) \setminus \{v\} \subseteq N(v) \setminus \{u\}$ ). Using this operation, one can simplify the decomposition theorem of Hermelin et al. [21], and thus we think this observation has an impact beyond this work.

Using the structural theorem, DISCONNECTED CUT on claw-free graphs reduces to understanding its behavior under the elementary operations and on the basic graph classes. The crucial elementary operation is to remove certain cobipartite

structures known as  $W$ -joins [6]. We develop the notion of unshatterable proper  $W$ -joins, which are essentially  $W$ -joins that cannot be broken into smaller  $W$ -joins, and exhibit how unshatterable proper  $W$ -joins interact with disconnected cuts. We then show that unshatterable proper  $W$ -joins can be removed from the graph by a simple operation. We complete our arguments by proving that all  $W$ -joins in the graph must in fact be unshatterable proper  $W$ -joins, and that we can find unshatterable proper  $W$ -joins in polynomial time.

The main basic graph classes in the structural theorem are line graphs and proper circular-arc graphs. Prior to our work, the complexity of DISCONNECTED CUT was unknown for these classes as well. We present a polynomial-time algorithm for line graphs and even for general circular-arcs graphs (not only proper circular-arcs). Both algorithms rely on the existence of a small induced cycle passing through a disconnected cut in a highly structured manner. In addition, for line graphs, we prove that the pre-image of the line graph is  $2P_2$ -free, and thus has diameter at most 3. The hardest part of the proof is then to prove that if the pre-image has diameter exactly 3, then the line graph has no disconnected cut.

## 1.2. Related work

As mentioned, the name DISCONNECTED CUT stems from Fleischner et al. [16], who studied how to partition the vertices of a graph into exactly  $k$  bicliques, where DISCONNECTED CUT is equivalent to the case  $k = 2$  in the complement graph. However, DISCONNECTED CUT originates from  $H$ -partitions, introduced in [9]. A model graph  $H$  on vertices  $h_1, \dots, h_k$  has solid and dotted edges. An  $H$ -partition of a graph  $G$  is a partition of  $V(G)$  into  $|V(H)|$  nonempty sets  $V_1, \dots, V_k$  such that for every pair of vertices  $u \in V_i$  and  $v \in V_j$ : if  $h_i h_j$  is a solid edge of  $E(H)$ , then  $uv \in E(G)$ ; and if  $h_i h_j$  is a dotted edge of  $E(H)$ , then  $uv \notin E(G)$  (if  $h_i h_j \notin E(H)$ , then  $uv \in E(G)$  or  $uv \notin E(G)$  are both allowed). The corresponding decision problem is called  $H$ -PARTITION. Dantas et al. [9] proved that  $H$ -PARTITION is polynomial-time solvable for every 4-vertex model graph  $H$  except  $H = 2K_2$ , which has solid edges  $h_1 h_3, h_2 h_4$  and no dotted edges, and  $H = 2S_2$ , which has dotted edges  $h_1 h_3, h_2 h_4$  and no solid edges. As a graph has a disconnected cut if and only if it has a  $2S_2$ -partition if and only if its complement has a  $2K_2$ -partition, these two cases are polynomial-time equivalent to DISCONNECTED CUT. Hence, we now know that, as a matter of exception,  $H$ -PARTITION is NP-complete if  $H \in \{2K_2, 2S_2\}$  [28].

We can encode a model graph  $H$  as a matrix  $M$  in which every entry is either 0 (dotted edge), 1 (solid edge) or \* (no restriction). If we allow sets  $V_i$  in a solution for  $H$ -PARTITION to be empty, then we obtain the  $M$ -PARTITION problem, introduced by Feder et al. [14]. This well-known problem generalizes many classical problems involving vertex cuts and partitions, including  $k$ -COLOURING and  $H$ -COLOURING; see also [19]. An even more general variant is to give every vertex  $u$  a list  $L(u) \subseteq \{1, \dots, k\}$  and to search for a solution, in which each vertex  $u$  may only belong to a set  $V_i$  with  $i \in L(u)$ . This yields the LIST  $M$ -PARTITION problem, which includes well-known cases, such as the STUBBORN problem, which turned out to be polynomial-time solvable [8], in contrast to DISCONNECTED CUT. A homomorphism  $f$  from  $G$  to  $H$  is a retraction if  $G$  contains  $H$  as an induced subgraph and  $f(u) = u$  for every  $u \in V(H)$ . The corresponding decision version is called  $H$ -RETRACTION. Let  $C_4$  be the 4-cycle with a self-loop in each vertex. Then  $C_4$ -RETRACTION is a special case of LIST  $2S_2$ -PARTITION where the input graph contains a cycle on four specified vertices  $v_1, \dots, v_4$  with  $L(v_i) = \{i\}$  for  $i = 1, \dots, 4$  and  $L(v) = \{1, 2, 3, 4\}$  for  $v \notin \{v_1, \dots, v_4\}$ . This problem is a generalization of DISCONNECTED CUT. Feder and Hell [13] proved that  $C_4$ -RETRACTION is NP-complete. Hence, LIST  $2S_2$ -PARTITION and LIST  $2K_2$ -PARTITION are NP-complete. Note that this result is also implied by the NP-completeness of  $2K_2$ -PARTITION [28].

Vikas [33] solved an open problem of Winkler (see [14,33]) by proving NP-completeness of  $C_4$ -COMPACTION, the variant of the  $2S_2$ -PARTITION problem with the extra constraint that there must be at least one edge  $u_i u_j$  with  $u_i \in V_i$  and  $u_{i+1} \in V_{i+1}$  for  $i = 1, \dots, 4$  (where  $V_5 = V_1$ ). Generally, a homomorphism  $f$  from a graph  $G$  to a graph  $H$  is a compaction if  $f$  is edge-surjective, i.e., for every  $xy \in E(H)$  with  $x \neq y$  there is an edge  $uv \in E(G)$  with  $f(u) = x$  and  $f(v) = y$ . The corresponding decision problem is called  $H$ -COMPACTION. If  $H = C_4$ , then the problem is equivalent to DISCONNECTED CUT when restricted to graphs of diameter 2 [16]. Hence,  $C_4$ -COMPACTION is NP-complete for graphs of diameter 2 [28] (the result of [33] holds for graphs of diameter at least 3). Similarly, a homomorphism  $f$  from a graph  $G$  to a graph  $H$  is (vertex-)surjective if for every  $x \in V(H)$  there is a vertex  $u \in V(G)$  such that  $f(u) = x$ . The decision problem is called SURJECTIVE  $H$ -COLOURING (or  $H$ -VERTEX COMPACTION, or SURJECTIVE  $H$ -HOMOMORPHISM) and is equivalent to DISCONNECTED CUT if  $H = C_4$ . The complexity classifications of  $H$ -COMPACTION and SURJECTIVE  $H$ -COLOURING are wide open despite many partial results; see [2] for a survey and [17] for a more recent overview focussing on SURJECTIVE  $H$ -COLOURING.

## 1.3. Overview

In Section 2 we state several underlying structural observations for graphs of diameter 2. In Sections 3 and 4, we prove that DISCONNECTED CUT can be solved in polynomial time for circular-arc graphs and line graphs, respectively. In Section 5 we prove our main result, and in particular, our new structural theorem for claw-free graphs of diameter 2. In Section 6 we show that DISCONNECTED CUT is polynomial-time solvable on paw-free graphs, co-paw-free graphs and on diamond-free graphs. By combining these results with our result for claw-free graphs we prove that DISCONNECTED CUT is polynomial-time solvable for  $H$ -free graphs whenever  $H$  is a graph on at most four vertices not isomorphic to the complete graph  $K_4$ . We pose the case where  $H = K_4$  as an open problem in Section 7, together with some other relevant open problems.

## 2. Preliminaries and basic results

In the remainder of our paper, graphs are finite, undirected, and have neither multiple edges nor self-loops unless explicitly stated otherwise.

Let  $G = (V, E)$  be a graph. For a set  $S \subseteq V$ ,  $G[S]$  is the subgraph of  $G$  induced by  $S$ . We say that  $S$  is *connected* if  $G[S]$  is connected. We write  $G - S = G[V \setminus S]$ , and if  $S = \{u\}$ , we write  $G - u$  instead. For a vertex  $u \in V$ , let  $N(u) = \{v \mid uv \in E\}$  be the neighbourhood of  $u$  and  $N[u] = N(u) \cup \{u\}$ . The *complement*  $\overline{G}$  of  $G$  has vertex set  $V$  and edge set  $\{uv \mid uv \notin E\}$ . The *contraction* of an edge  $uv \in E$  is the operation that removes the vertices  $u$  and  $v$  from  $G$ , and replaces  $u$  and  $v$  by a new vertex that is made adjacent to precisely those vertices that were adjacent to  $u$  or  $v$  in  $G$  (without introducing self-loops or multiple edges). The *distance*  $d_G(u, v)$  between two vertices  $u$  and  $v$  of  $G$  is the number of edges in a shortest path between them. If  $u$  and  $v$  are in different connected components of  $G$ , then  $d_G(u, v) = \infty$ . The *diameter* of  $G$  is equal to  $\max\{d_G(u, v) \mid u, v \in V\}$ .

The following lemma was observed by Fleischner et al.

**Lemma 1** ([16]). *Let  $G$  be a graph. If  $G$  has diameter 1, then  $G$  has no disconnected cut. If  $G$  has diameter at least 3, then  $G$  has a disconnected cut, which can be found in linear time.*

A subset  $D \subseteq V$  is a *dominating set* of a graph  $G = (V, E)$  if every vertex of  $V \setminus D$  is adjacent to at least one vertex of  $D$ . If  $D = \{u\}$ , then  $u$  is a *dominating vertex* of  $G$ . An edge  $uv$  of a graph  $G = (V, E)$  is *dominating* if  $\{u, v\}$  is dominating. A vertex  $u \in V$  has a *disconnected neighbourhood* if  $N(u)$  induces a disconnected graph.

We need the following two lemmas, the first one of which is a straightforward observation.

**Lemma 2.** *If a graph  $G$  contains a dominating vertex, then  $G$  has no disconnected cut.*

**Lemma 3.** *If a graph  $G$  contains a non-dominating vertex  $u$  with a disconnected neighbourhood, then  $G$  has a disconnected cut.*

**Proof.** Let  $A_1, \dots, A_r$  be the connected components of  $G[N(u)]$  for some  $r \geq 2$ . As  $u$  is not dominating,  $G - (N(u) \cup \{u\})$  is nonempty. We define  $V_1 = \{u\}$ ,  $V_2 = V(A_1)$ ,  $V_3 = V(G) - (N(u) \cup \{u\})$ , and  $V_4 = V(A_2) \cup \dots \cup V(A_r)$ , and note that  $V_1 \cup V_3$  (or  $V_2 \cup V_4$ ) is a disconnected cut of  $G$ .  $\square$

Two disjoint vertex sets  $S$  and  $T$  in a graph  $G = (V, E)$  are *complete* to each other if there is an edge between every vertex of  $S$  and every vertex of  $T$ , and  $S$  and  $T$  are *anticomplete* to each other if there is no edge between a vertex of  $S$  and a vertex of  $T$ . Recall that  $G$  has a *disconnected cut* if  $V$  can be partitioned into four (nonempty) sets  $V_1, V_2, V_3, V_4$ , such that  $V_1$  is anticomplete to  $V_3$  and  $V_2$  is anticomplete to  $V_4$ . We say that  $V_1, V_2, V_3, V_4$  form a *disconnected partition* of  $G$ .

**Lemma 4.** *Let  $V_1, V_2, V_3, V_4$  be a disconnected partition of a graph  $G$  of diameter 2. Then  $G$  has an induced cycle  $C$  with  $4 \leq |V(C)| \leq 5$  such that  $V(C) \cap V_i \neq \emptyset$  for  $i = 1, \dots, 4$ .*

**Proof.** Let  $u_1 \in V_1$  and  $u_3 \in V_3$ . As  $G$  has diameter 2, there exists a vertex  $u_2$  in  $V_2$  or  $V_4$ , say  $V_2$ , such that  $u_2$  is adjacent to  $u_1$  and to  $u_3$ . Let  $u_4 \in V_4$ . As  $G$  has diameter 2, there exists a vertex  $u'_1$  in  $V_1$  or  $V_3$ , say  $V_1$ , such that  $u'_1$  is adjacent to  $u_2$  and  $u_4$ . If  $u_3$  and  $u_4$  are adjacent, then we can take as  $C$  the cycle on vertices  $u'_1, u_2, u_3, u_4$  in that order. Otherwise, as  $G$  has diameter 2, there exists a vertex  $w \in V_3 \cup V_4$ , such that  $w$  is adjacent to  $u_3$  and to  $u_4$ . In that case we can take as  $C$  the cycle on vertices  $u'_1, u_2, u_3, w, u_4$ .  $\square$

Two adjacent vertices  $u$  and  $v$  of graph  $G = (V, E)$  have a *nested neighbourhood* if  $N(u) \setminus \{v\} \subseteq N(v) \setminus \{u\}$  or  $N(v) \setminus \{u\} \subseteq N(u) \setminus \{v\}$ . We say that  $G$  has *distinct neighbourhoods* if  $G$  has no two vertices that have nested neighbourhoods. In our proof we will apply the following lemma exhaustively.

**Lemma 5.** *Let  $G$  be a graph of diameter 2 that contains two vertices  $u$  and  $v$  such that  $N(u) \setminus \{v\} \subseteq N(v) \setminus \{u\}$ . Then  $G$  has a disconnected cut if and only if  $G - u$  has a disconnected cut. Moreover,  $G - u$  has diameter at most 2.*

**Proof.** As  $G$  has diameter 2 and  $N(u) \setminus \{v\} \subseteq N(v) \setminus \{u\}$ , we find that  $G - u$  has diameter at most 2.

First suppose that  $G$  has a disconnected cut. Then  $G$  has a disconnected partition  $V_1, V_2, V_3, V_4$ . We may assume without loss of generality that  $v$  belongs to  $V_1$ . First assume that  $u \in V_1$ . Hence,  $\{u, v\} \subseteq V_1$  and thus,  $V_1 \setminus \{u\}, V_2, V_3, V_4$  form a disconnected partition of  $G - u$ . Hence,  $G - u$  has a disconnected cut. Now assume that  $u \notin V_1$ . As  $u$  is adjacent to  $v$ , we may assume without loss of generality that  $u$  belongs to  $V_2$ . As  $v$  belongs to  $V_1$  and  $V_1$  is anticomplete to  $V_3$ , we find that  $v$  has no neighbours in  $V_3$ . As  $N(u) \setminus \{v\} \subseteq N(v) \setminus \{u\}$ , this means that  $u$  has no neighbours in  $V_3$ . As  $G$  has diameter 2, this means that  $V_2$  contains a vertex  $w \neq u$  that has a neighbour in  $V_3$ , thus  $V_2 \setminus \{u\}$  is nonempty. As a consequence,  $V_1, V_2 \setminus \{u\}, V_3, V_4$  form a disconnected partition of  $G - u$ . Hence,  $G - u$  has a disconnected cut.

Now suppose that  $G - u$  has a disconnected cut. Then  $G - u$  has a disconnected partition  $V'_1, V'_2, V'_3, V'_4$ . We may assume without loss of generality that  $v$  belongs to  $V'_1$ . Then, as  $N(u) \setminus \{v\} \subseteq N(v) \setminus \{u\}$ , we find that  $V'_1 \cup \{u\}, V'_2, V'_3, V'_4$  form a disconnected partition of  $G$ . Hence,  $G$  has a disconnected cut.  $\square$

A pair of vertices  $u$  and  $v$  of a graph  $G = (V, E)$  is a *universal pair* if  $\{u, v\}$  is a dominating set and there exist distinct vertices  $x$  and  $y$  in  $V \setminus \{u, v\}$ , such that  $x \in N(u)$  and  $y \in N(v)$ ; note that this implies that  $|V| \geq 4$  and  $u, v$  have at least one neighbour in  $V \setminus \{u, v\}$ . Let  $H$  be a graph. Then  $G$  is  *$H$ -free* if  $G$  contains no induced subgraph isomorphic to  $H$ . The *disjoint union*  $G + H$  of two vertex-disjoint graphs  $G$  and  $H$  is the graph  $(V(G) \cup V(H), E(G) \cup E(H))$ . The disjoint union of  $r$  copies of a graph  $G$  is denoted by  $rG$ . The graphs  $C_r$  and  $P_r$  denote the cycle and path on  $r$  vertices, respectively. The graph  $K_r$  denotes the complete graph on  $r$  vertices. The *independence number*  $\alpha(G)$  of a graph  $G$  is the largest  $k$  such that  $G$  contains an induced subgraph isomorphic to  $kP_1$ .

**Lemma 6** ([7]). *A  $2P_2$ -free graph has a disconnected cut if and only if its complement has a universal pair.*

**Lemma 7** ([10]). *DISCONNECTED CUT is  $O(n^3)$ -time solvable for  $4P_1$ -free graphs.*

A graph  $G$  is *bipartite* if  $V(G)$  can be partitioned into two classes  $A$  and  $B$  such that every edge of  $G$  has an endpoint in  $A$  and an endpoint in  $B$ . If  $A$  is complete to  $B$ , then  $G$  is a *complete bipartite graph*. The graph  $K_{s,t}$  denotes the complete bipartite graph with partition classes of size  $s$  and  $t$ , respectively. The graph  $(\{u, v_1, v_2, v_3\}, \{uv_1, uv_2, uv_3\})$  is the *claw*  $K_{1,3}$ . A graph is *cobipartite* if it is the complement of a bipartite graph.

The *line graph* of a graph  $G$  with edges  $e_1, \dots, e_p$  is the graph  $L(G)$  with vertices  $u_1, \dots, u_p$  such that there is an edge between any two vertices  $u_i$  and  $u_j$  if and only if  $e_i$  and  $e_j$  have a common endpoint in  $G$ . Note that every line graph is claw-free. We call  $G$  the *preimage* of  $L(G)$ . Every connected line graph except  $K_3$  has a unique preimage [18].

A *circular-arc graph* is a graph that has a representation in which each vertex corresponds to an arc of a circle, such that two vertices are adjacent if and only if their corresponding arcs intersect. An *interval graph* is a graph that has representation in which each vertex corresponds to an interval of the line, such that two vertices are adjacent if and only if their corresponding intervals intersect. Note that circular-arc graphs generalize interval graphs. A circular-arc or interval graph is *proper* if it has a representation where the arcs respectively intervals are such that no one is contained in another. A *chordal graph* is a graph in which every induced cycle is a triangle; note that every interval graph is chordal, but that there exist circular-arc graphs, such as cycles, that are not chordal.

### 3. Circular-arc graphs

In this section we prove that DISCONNECTED CUT is polynomial-time solvable for circular-arc graphs. This result is known already for interval graphs, as it follows from the result that DISCONNECTED CUT is polynomial-time solvable for the class of chordal graphs [23], which contains the class of interval graphs. In fact, we have an  $O(n^2)$ -time algorithm for interval graphs. Due to Lemma 4 and the fact that interval graphs are chordal, no interval graph of diameter 2 has a disconnected cut. Consequently, an interval graph has a disconnected cut if and only if its diameter is at least 3 due to Lemma 1. To show that DISCONNECTED CUT is polynomial-time solvable for circular-arc graphs requires significant additional work.

Let  $G$  be a circular-arc graph. For each vertex  $u \in V(G)$  we can associate an arc  $[l_u, r_u]$  where we say that  $l_u$  is the clockwise left endpoint of  $u$  and  $r_u$  is the clockwise right endpoint of  $u$ . The following result of McConnell shows that we may assume that all left and right endpoints of the vertices of  $G$  are unique.

**Lemma 8** ([30]). *A circular-arc graph  $G$  on  $n$  vertices and  $m$  edges can be recognized in  $O(n + m)$  time. In the same time, a representation of  $G$  can be constructed with distinct arc endpoints that are clockwise enumerated as  $1, \dots, 2n$ .*

We need the following lemma.

**Lemma 9.** *Let  $G$  be a circular-arc graph of diameter 2 with a disconnected cut. Then  $G$  has a disconnected partition  $V_1, V_2, V_3, V_4$  such that each  $V_i$  is connected.*

**Proof.** By Lemma 8 we may assume that  $G$  has a representation with distinct arc endpoints clockwise enumerated as  $1, \dots, 2n$ . Let  $V_1, V_2, V_3, V_4$  be a disconnected partition of  $G$ . By Lemma 4,  $G$  contains a cycle  $C$  with vertices  $u_i$  for  $i = 1, \dots, j$  (with  $u_{j+1} = u_1$ ) and  $j \in \{4, 5\}$ , such that  $V(C) \cap V_i \neq \emptyset$  for  $i = 1, 2, 3, 4$ . We may assume without loss of generality that if  $j = 5$ , then  $u_5 \in V_4$ .

Let  $D_i$  be the connected component of  $G[V_i]$  that contains  $u_i$  for  $i = 1, \dots, 4$ . Note that if  $j = 5$ , then  $u_5 \in D_4$ . We let  $[l_i, r_i]$  denote the arc corresponding to the arc covered by the vertices of  $D_i$  (or equivalently, the arc associated with the vertex obtained after contracting the connected graph  $D_i$  into a single vertex). Note that the arcs  $[l_1, r_1], \dots, [l_4, r_4]$  cover the whole circle, as they contain the arcs corresponding to the vertices of  $C$ . Moreover,  $[l_i, r_i]$  intersects with  $[l_{i-1}, r_{i-1}]$  and  $[l_{i+1}, r_{i+1}]$  for  $i = 1, \dots, 4$  (where  $[l_0, r_0] = [l_5, r_5] = [l_1, r_1]$ ).

We set  $V'_i := V(D_i)$ . If  $V'_i = V_i$  for  $i = 1, \dots, 4$ , then  $V_1, V_2, V_3, V_4$  is our desired partition of  $G$ . Assume that this is not the case. Then  $V'_1 \cup \dots \cup V'_4 \subsetneq V$ . Consider an arbitrary vertex  $v \in V(G)$  that is not in any  $V'_i$ . Let  $p$  be the number of arcs  $[l_i, r_i]$  that intersect with  $[l_v, r_v]$ . We claim that  $p \leq 1$ .

First suppose that  $p = 4$ . Then  $v$  would be adjacent to a vertex of every  $V_i$ . This is not possible, because  $V_1, V_2, V_3, V_4$  is a disconnected partition of  $G$ . Now suppose that  $p = 3$ , say  $[l_v, r_v]$  intersects with  $[l_1, r_1], [l_2, r_2]$  and  $[l_3, r_3]$ . Then  $v$  must be in  $V_2$ , that is,  $v$  is in the same set as the vertices of  $D_2$ . As  $v$  intersects with  $[l_2, r_2]$  we would have put  $v$  in  $D_2$ , a contradiction. Now suppose that  $p = 2$ . By construction,  $[l_v, r_v]$  intersects with two consecutive arcs, say with arcs  $[l_1, r_1]$  and  $[l_2, r_2]$ . Then  $v \in V_1$  or  $v \in V_2$ . If  $v \in V_1$  then we would have put  $v$  in  $D_1$ , and if  $v \in V_2$  then we would have put  $v \in D_2$ . Hence, this situation is not possible either.

From the above, we conclude that  $p \leq 1$ , that is, for each  $v \notin V'_1 \cup \dots \cup V'_4$ , we have that  $[l_v, r_v]$  intersects with at most one  $[l_i, r_i]$ . As the arcs  $[l_1, r_1], \dots, [l_4, r_4]$  cover the whole circle, this means that  $[l_v, r_v]$  is contained in some  $[l_i, r_i]$ , and we can safely put  $v$  in  $V'_i$  without destroying the connectivity or the anticompleteness of  $V_1, V_3$ , or of  $V_2, V_4$ . Afterwards, the four sets  $V'_1, V'_2, V'_3, V'_4$  form a disconnected partition of  $G$ , such that each  $V'_i$  is connected.  $\square$

We are now ready to prove the main result of this section.

**Theorem 1.** DISCONNECTED CUT is  $O(n^2)$ -time solvable for circular-arc graphs.

**Proof.** Let  $G = (V, E)$  be a circular-arc graph on  $n$  vertices. We will give an  $O(n^2)$ -time algorithm that either finds a disconnected cut of  $G$  or concludes that  $G$  has no disconnected cut.

We compute the diameter of  $G$  in  $O(n^2)$  time using the (more general)  $O(n^2)$ -time algorithm of [5] (or of [1] or the linear-time algorithm of [27]). Lemma 1 tells us that if  $G$  has diameter 1, then  $G$  has no disconnected cut, and if  $G$  has diameter at least 3, then  $G$  has a disconnected cut. Assume that  $G$  has diameter 2. Lemma 9 tells us that if  $G$  has a disconnected cut, then  $G$  has a disconnected partition  $V_1, V_2, V_3, V_4$  such that  $V_i$  is connected for  $i = 1, \dots, 4$ . We say that the arc of a set  $V_i$  is the union of all the arcs of the vertices in  $V_i$ . As  $G$  has diameter 2, the union of the arcs of the sets  $V_i$  covers the whole circle. Moreover, the arcs of  $V_1$  and  $V_3$  are disjoint and the arcs of  $V_2$  and  $V_4$  are disjoint.

We now compute, in linear time, a representation of  $G$  with distinct arc endpoints clockwise enumerated as  $1, \dots, 2n$  via Lemma 8. After sorting the arcs in  $O(n \log n)$  time, we apply the following procedure for each  $v \in V$ . We take a neighbour  $v'$  with right-most right endpoint amongst all neighbours of  $v$ . We then take a neighbour  $v''$  with right-most right endpoint amongst all neighbours of  $v'$ . We check if  $v''$  is also a neighbour of  $v$ . If so, then  $v, v', v''$  form a triangle if  $v \neq v''$  and an edge if  $v = v''$  such that the corresponding arcs cover the whole circle. In that case  $G$  has no disconnected partition  $V_1, V_2, V_3, V_4$  such that  $V_i$  is connected for  $i = 1, \dots, 4$  (as the arcs of the three vertices of the triangle must be placed in the corresponding arcs of the sets  $V_i$ ), and then, by Lemma 9, we find that  $G$  has no disconnected cut. Note that this procedure takes  $O(n^2)$  time in total.

Suppose that we did not find a pair or triple of vertices in  $G$  whose arcs cover the whole circle. We now aim to find a small induced cycle  $v_1, \dots, v_k$  as follows. Let  $v_1$  be an arbitrary vertex of  $G$  whose arc is not contained in the arc of another vertex. Now iteratively perform the following procedure starting with  $v_i$ : let  $v_{i+1}$  be the neighbour of  $v_i$  with the right-most right endpoint amongst all neighbours of  $v_i$ . We continue until  $v_{i+1}$  neighbours  $v_1$ . The procedure runs in  $O(n)$  time and yields an induced cycle  $v_1, \dots, v_k$ , unless the procedure gets stuck without finding a vertex neighbouring  $v_1$ . In that case,  $G$  has no induced cycle on four or more vertices, and thus no disconnected cut due to Lemma 4. So assume that we do find an induced cycle. As  $G$  has diameter 2 and  $G$  has no pair or triple of vertices whose arcs cover the whole circle, we find that  $k \in \{4, 5\}$ . Moreover, by construction and because  $G$  is circular-arc, the arcs corresponding to the vertices  $v_i$  must cover the whole circle.

If  $G$  has a disconnected partition  $V_1, V_2, V_3, V_4$  such that each  $V_i$  is connected, then the above implies the following. If  $k = 4$ , then we may assume without loss of generality that  $v_i \in V_i$  for  $i = 1, \dots, 4$ . If  $k = 5$ , then two vertices  $v_i, v_{i+1}$  belong to the same set  $V_h$ , whereas sets  $V_i$  with  $i \neq h$  each contain a single vertex from  $C$ . If  $k = 5$ , then we guess which two vertices  $v_i, v_{i+1}$  will be put in the same set, say  $v_1, v_5$ ; this does not influence the asymptotic running time.

Now we build up the sets  $V_i$  from scratch by putting in the vertices from  $V(G) \setminus \{v_1, \dots, v_k\}$ . We will always maintain that each  $V_i$  induces a connected graph, and thus, the union of the arcs of the vertices in  $V_i$  indeed always form an arc. Observe also that no set  $V_h$  is contained in some other set  $V_i$ ; by our choice of vertices  $v_i$  we will always maintain this property. We say that a vertex  $u$  intersects a set  $V_i$  if the arc of  $u$  intersects the arc of  $V_i$ . Note that, since the arcs corresponding to  $\{v_1, \dots, v_k\}$  cover the entire circle, so do the arcs of the sets  $V_i$  that we are constructing. If there is a vertex that intersects each of the sets  $V_i$  constructed so far, then there is no disconnected cut with each  $V_i$  connected. If  $k = 4$ , then  $G$  has no disconnected cut due to Lemma 9. If  $k = 5$ , then our guess of vertices  $v_1, v_5$  to belong to  $V_1$  may have been incorrect, and we need to put two other consecutive vertices of  $C$  in the same set  $V_h$  before concluding that  $G$  has no disconnected cut. Otherwise, we do the following until no longer possible. As no  $V_h$  is contained in some other  $V_h$ , any vertex  $u$  that intersects two sets  $V_i$  and  $V_{i+2}$  for some  $i$  (say,  $i \in \{1, 2\}$  without loss of generality), also intersects  $V_{i+1}$  or  $V_{i+3}$  (where  $V_5 = V_1$ ). We now put a vertex  $u$  that intersects two sets  $V_i$  and  $V_{i+2}$  for some  $i$  into set  $V_{i+1}$  if  $u$  intersects  $V_{i+1}$  as well; otherwise,  $u$  intersects  $V_{i+3}$  and we put  $u$  in  $V_{i+3}$ .

Let  $T$  be the set of vertices of  $G$  that we have not placed in some set  $V_i$  yet. We claim that each vertex of  $T$  must intersect exactly two sets  $V_i$  and  $V_j$  such that, in addition,  $j = i + 1$  holds. For contradiction, assume that this does not

hold for  $u \in T$ . Then  $u$  intersects exactly one set  $V_i$ , say  $V_1$ . There must exist a path from  $u$  to  $v_3$  of length 2, as  $G$  has diameter 2. Hence, there exists a vertex  $w$  that is adjacent to both  $u$  and  $v_3$ . This means that the arc corresponding to  $w$  must intersect  $V_1$  and  $V_3$ . Hence,  $w$  has already been placed in  $V_2$  or  $V_4$  and thus the arc of  $u$  intersects two sets, namely  $V_1$  and  $V_2$ , or  $V_1$  and  $V_4$ , a contradiction. The claim follows.

As every vertex in  $T$  intersects two sets  $V_i$  and  $V_{i+1}$ , we can model the remaining instance as an instance of 2-SATISFIABILITY as follows. Let  $u \in T$ , where  $u$  must be placed in, say,  $V_i$  or  $V_{i+1}$ . We introduce two variables  $x_u^i$  and  $x_u^{i+1}$  with clauses  $x_u^i \vee x_u^{i+1}$  and  $\bar{x}_u^i \vee \bar{x}_u^{i+1}$ . For each edge  $uv$  where  $u$  must be placed in  $V_i$  or  $V_{i+1}$  and  $v$  must be placed in  $V_{i+1}$  or  $V_{i+2}$ , we introduce the clause  $\bar{x}_u^i \vee \bar{x}_v^{i+2}$ . For each edge  $uv$  where  $u$  must be placed in  $V_i$  or  $V_{i+1}$  and  $v$  must be placed in  $V_{i+2}$  or  $V_{i+3}$ , we introduce the clauses  $\bar{x}_u^i \vee \bar{x}_v^{i+2}$  and  $\bar{x}_u^{i+1} \vee \bar{x}_v^{i+3}$ .

The correctness of our algorithm follows from the above description. Assigning the vertices to the sets  $V_i$  takes  $O(n^2)$  time, whereas solving the corresponding instance of 2-SATISFIABILITY takes  $O(n^2)$  time as well. As computing the diameter takes  $O(n^2)$  time and all other steps take  $O(n^2)$  time as well, the total running time is  $O(n^2)$ .  $\square$

#### 4. Line graphs

In this section we prove that DISCONNECTED CUT is polynomial-time solvable for line graphs. We start with the following lemma due to Ito et al. [23].

**Lemma 10 ([23]).** *Let  $G$  be a graph with diameter 2 whose line graph  $L(G)$  also has diameter 2. Then  $G$  has a disconnected cut if and only if  $L(G)$  has a disconnected cut.*

We need a lemma on graphs whose line graph has diameter 2.

**Lemma 11.** *Let  $G$  be a graph that is neither a triangle nor a star. Then  $L(G)$  has diameter 2 if and only if  $G$  is  $2P_2$ -free.*

**Proof.** Let  $G$  be a graph that is neither a triangle nor a star. First suppose that  $L(G)$  has diameter 2. For contradiction, assume that  $G$  is not  $2P_2$ -free. Then  $G$  contains an induced subgraph  $H$  with four distinct vertices  $s, t, u, v$  and edges  $e_1 = st$  and  $e_2 = uv$ . As  $H$  is an induced subgraph of  $G$ , we find that  $e_1$  and  $e_2$  are non-adjacent vertices in  $L(G)$ . Then, because  $L(G)$  has diameter 2,  $L(G)$  contains a vertex  $e_3$  that is adjacent to  $e_1$  and  $e_2$ . However, then  $e_3$  is an edge with one endpoint in  $\{s, t\}$  and the other one in  $\{u, v\}$ . This means that  $H$  is not induced, a contradiction.

Now suppose that  $G$  is  $2P_2$ -free. For contradiction, assume that  $L(G)$  has diameter not equal to 2. If the diameter of  $L(G)$  is 1, then  $L(G)$  is a complete graph implying that  $G$  is a triangle or star, which is not what we assume. If the diameter of  $L(G)$  is at least 3, then  $L(G)$  contains two vertices  $e_1$  and  $e_2$  that are of distance at least 3. This means that  $e_1$  and  $e_2$  form an induced  $2P_2$  in  $G$ , a contradiction.  $\square$

We are now ready to prove the main result of this section.

**Theorem 2.** *The DISCONNECTED CUT problem is  $O(n^4)$ -time solvable for line graphs of  $n$ -vertex graphs.*

**Proof.** Let  $G$  be a graph on  $n$  vertices and  $m$  edges. We will either find in  $O(n^4)$  time that  $L(G)$  has a disconnected cut or conclude that  $L(G)$  has no disconnected cut. We first check in  $O(n)$  time if  $G$  is a triangle or star. If so, then  $L(G)$  is a complete graph and thus  $L(G)$  has no disconnected cut. From now on suppose that  $G$  is neither a triangle nor a star. By Lemma 11 we find that  $L(G)$  has diameter 2 if and only if  $G$  is  $2P_2$ -free. Hence, we can check in  $O(n^4)$  time, via checking if  $G$  has an induced  $2P_2$  by brute force, if  $L(G)$  has diameter 2.

First assume that  $L(G)$  does not have diameter 2. As  $G$  is not a triangle or a star,  $L(G)$  has diameter at least 3. By Lemma 1 we find that  $L(G)$  has a disconnected cut. Now assume that  $L(G)$  has diameter 2. We check in  $O(n^3)$  time if  $G$  has an edge  $uv$  such that every vertex of  $V(G) \setminus \{u, v\}$  is adjacent to at least one of  $u, v$ . If so, then  $uv$  is a dominating vertex of  $L(G)$ , and  $L(G)$  has no disconnected cut due to Lemma 2. If not, then  $L(G)$  has no dominating vertices, and we proceed as follows. First we check if  $L(G)$  has a vertex  $uv$  with a disconnected neighbourhood, or equivalently, if  $G$  contains an edge  $uv$  such that  $u$  and  $v$  have degree at least 2 and no common neighbours. This takes  $O(n^3)$  time. If  $L(G)$  has a vertex with a disconnected neighbourhood, then  $L(G)$  has a disconnected cut by Lemma 3. From now on assume that  $L(G)$  has no vertex with a disconnected neighbourhood. As  $G$  is neither a triangle nor a star,  $G$  is  $2P_2$ -free by Lemma 11. Hence,  $G$  has diameter at most 3. We can determine in  $O(n^3)$  time the diameter of  $G$  and consider each case separately.

**Case 1.**  $G$  has diameter 1.

We claim that  $L(G)$  has no disconnected cut. For contradiction, assume that  $L(G)$  has a disconnected cut. Let  $V'_1, V'_2, V'_3, V'_4$  be a disconnected partition of  $L(G)$ . By Lemma 4,  $L(G)$  contains a cycle  $C'$  with vertices  $u_i u_{i+1}$  for  $i = 1, \dots, j$  (with  $u_{j+1} = u_1$ ) and  $j \in \{4, 5\}$ , such that  $V(C') \cap V'_i \neq \emptyset$  for  $i = 1, 2, 3, 4$ . Then we may assume without loss of generality that  $u_i u_{i+1} \in V'_i$  for  $i = 1, \dots, 4$  and  $u_j u_{j+1} \in V'_4$ . As  $G$  has diameter 1,  $u_1 u_3$  is an edge of  $G$  and thus a vertex of  $L(G)$ . In  $L(G)$ ,  $u_1 u_3$  is adjacent to every vertex in  $\{u_1 u_2, u_2 u_3, u_3 u_4, u_j u_{j+1}\}$ , and thus to a vertex in  $V'_i$  for  $i = 1, \dots, 4$ , a contradiction.

**Case 2.**  $G$  has diameter 2.

Then  $G$  has a disconnected cut if and only if  $L(G)$  has a disconnected cut due to Lemma 10. By Lemma 6 it suffices to check if  $\overline{G}$  has a universal pair. This takes  $O(n^3)$  time.

**Case 3.**  $G$  has diameter 3.

We will prove that  $L(G)$  has no disconnected cut. As  $G$  has diameter 3,  $G$  does have a disconnected cut by Lemma 1. We need the following claim.

*Claim.* Let  $V_1, V_2, V_3, V_4$  be a disconnected partition of  $G$ . Then every cycle  $C$  of  $G$  with  $4 \leq |V(C)| \leq 5$  contains vertices of at most three distinct sets from  $\{V_1, V_2, V_3, V_4\}$ .

We prove the Claim as follows. For contradiction, assume that  $G$  has a cycle  $C$  with vertices  $u_1, \dots, u_j$  for  $j \in \{4, 5\}$ , such that  $V(C) \cap V_i \neq \emptyset$  for  $i = 1, \dots, 4$ . We may assume without loss of generality that  $u_i \in V_i$  for  $i = 1, \dots, 4$  and  $u_j \in V_4$ . As  $G$  is  $2P_2$ -free, we may assume without loss of generality that  $u_3$  is in a singleton connected component of  $G[V_3]$ . If  $j = 4$ , then we may also assume without loss of generality that  $u_2$  is in a singleton connected component of  $G[V_2]$ . If  $j = 5$ , then  $u_2$  must be in a singleton connected component of  $G[V_2]$  due to the edge  $u_4u_5$ , which is contained in  $G[V_4]$ . This means that the sets  $N_G(u_2) \setminus \{u_3\}$  and  $N_G(u_3) \setminus \{u_2\}$  are disjoint. As  $u_1u_2$  and  $u_3u_4$  are edges of  $G$ , both  $N_G(u_2) \setminus \{u_3\}$  and  $N_G(u_3) \setminus \{u_2\}$  are nonempty. Hence, the vertex  $u_2u_3$  has a disconnected neighbourhood in  $L(G)$ , a contradiction. This proves the Claim.

Now, for contradiction, assume that  $L(G)$  has a disconnected cut. Let  $V'_1, V'_2, V'_3, V'_4$  be a disconnected partition of  $L(G)$ . By Lemma 4,  $L(G)$  contains a cycle  $C'$  with vertices  $u_iu_{i+1}$  for  $i = 1, \dots, j$  (with  $u_{j+1} = u_1$ ) and  $j \in \{4, 5\}$ , such that  $V(C') \cap V'_i \neq \emptyset$  for  $i = 1, 2, 3, 4$ . Then we may assume without loss of generality that  $u_iu_{i+1} \in V'_i$  for  $i = 1, \dots, 4$  and  $u_ju_{j+1} \in V'_4$ .

We define the following partition  $V_1, V_2, V_3, V_4$  of  $V(G)$ . Let  $u \in V(G)$ . If  $u$  is incident to only edges from one set  $V'_i$ , then we put  $u$  in  $V_i$ . Suppose  $u$  is incident to edges from more than one set  $V'_i$ . As  $V'_1, V'_2, V'_3, V'_4$  is a disconnected partition of  $L(G)$ , we find that  $u$  is incident to edges from  $V'_i$  and  $V'_{i+1}$  for some  $1 \leq i \leq 4$  (where  $V'_5 = V'_1$ ) and to no other sets  $V'_j$ . In that case we put  $u$  into  $V_{i+1}$ .

We now prove that  $V_1$  is anticomplete to  $V_3$ . For contradiction, assume that  $V_1$  contains a vertex  $u$  and  $V_3$  contains a vertex  $v$  such that  $uv \in E(G)$ . As  $u \in V_1$ , we find that  $u$  is incident to edges only in  $V'_4$  and  $V'_1$ . Hence,  $uv \in V'_1 \cup V'_4$ . As  $v \in V_3$ , we find that  $v$  is incident to edges only in  $V'_2$  and  $V'_3$ . This implies that  $uv \in V'_2 \cup V'_3$ , a contradiction. By the same argument we can show that  $V_2$  is anticomplete to  $V_4$ . Let  $C$  be the cycle with vertices  $u_1, \dots, u_j$  in  $G$ . Then  $V(C) \cap V_i \neq \emptyset$ , and thus  $V_i \neq \emptyset$ , for  $i = 1, \dots, 4$ . Hence,  $V_1, V_2, V_3, V_4$  is a disconnected partition of  $G$ , and  $C$  is a cycle in  $G$  with  $V(C) \cap V_i \neq \emptyset$  for every  $i$ . This is not possible due to the Claim. We conclude that  $L(G)$  has no disconnected cut.

The correctness of our algorithm follows from the above description. If  $G$  has diameter 1 (Case 1) or diameter 3 (Case 3), no additional running time is required, as we showed that  $L(G)$  has no disconnected cut in both these cases. Hence, only executing Case 2 takes additional time, namely time  $O(n^3)$ . Hence, the total running time of our algorithm is  $O(n^4)$ , as desired.  $\square$

## 5. Claw-free graphs

In this section, we prove that DISCONNECTED CUT is polynomial-time solvable on claw-free graphs. The proof consists of two parts. In Section 5.1 we show how to get rid of certain cobipartite structures in the graph, called  $W$ -joins. We remark that DISCONNECTED CUT can be solved in polynomial time on cobipartite graphs [16]. Although this is a necessary condition for DISCONNECTED CUT to be solvable in polynomial time on claw-free graphs, the algorithm for cobipartite graphs is not sufficient to deal with  $W$ -joins. In Section 5.2 we present our new decomposition theorem for claw-free graphs of diameter 2 and combine this theorem with the results from the previous sections and Section 5.1 to show our main result.

### 5.1. Cobipartite structures versus disconnected cuts

We consider the following cobipartite structures that might be present in claw-free graphs [6,21]. A pair  $(A, B)$  of disjoint non-empty sets of vertices is a  $W$ -join in graph  $G$  if  $|A| + |B| > 2$ ,  $A$  and  $B$  are cliques,  $A$  is neither complete nor anticomplete to  $B$ , and every vertex of  $V(G) \setminus (A \cup B)$  is either complete or anticomplete to  $A$  and either complete or anticomplete to  $B$ . A  $W$ -join is a *proper*  $W$ -join if each vertex in  $A$  is neither complete nor anticomplete to  $B$  and each vertex in  $B$  is neither complete nor anticomplete to  $A$ . Observe that for a proper  $W$ -join  $(A, B)$ , it must hold that  $|A|, |B| \geq 2$ . For any  $W$ -join  $(A, B)$ , it holds that  $G[A \cup B]$  is a cobipartite induced subgraph in  $G$ .

We assume that an input graph  $G$  of DISCONNECTED CUT has diameter 2 and that  $G$  has distinct neighbourhoods, by Lemmas 1 and 5 respectively. We show how to use these assumptions to remove all  $W$ -joins in a claw-free graph and obtain an equivalent instance of DISCONNECTED CUT. As a first step, we show that we can focus on proper  $W$ -joins.

**Lemma 12.** Let  $G$  be a graph with distinct neighbourhoods. If  $G$  admits a  $W$ -join  $(A, B)$ , then  $(A, B)$  is a proper  $W$ -join.

**Proof.** We need to show that no vertex of  $A$  (respectively  $B$ ) is complete or anticomplete to  $B$  (respectively  $A$ ). Suppose there exists a vertex  $a \in A$  such that  $a$  is anticomplete to  $B$ . Since  $A$  is not anticomplete to  $B$ , there exists a vertex  $a' \in A \setminus \{a\}$



such that  $a'$  is adjacent to some vertex of  $B$ . This implies that  $N(a) \setminus \{a'\} \subseteq N(a') \setminus \{a\}$ , which is a contradiction to the fact that  $G$  has distinct neighbourhoods. Hence, no vertex of  $A$  is anticomplete to  $B$ . Similarly, no vertex of  $B$  is anticomplete to  $A$ .

Suppose there exists a vertex  $a \in A$  such that  $a$  is complete to  $B$ . Then for every  $a' \in A \setminus \{a\}$ , it holds that  $N(a') \setminus \{a\} \subseteq N(a) \setminus \{a'\}$ . Since  $G$  has distinct neighbourhoods, no such  $a'$  exists, and thus  $|A| = 1$ . Hence,  $|B| \geq 2$  by the definition of a  $W$ -join. However, no vertex of  $B$  is anticomplete to  $A$ . Since  $|A| = 1$ , this implies that each vertex of  $B$  is complete to  $A$ . Following the same reasoning as before, this implies that  $|B| = 1$ , a contradiction. Hence, no vertex of  $A$  is complete to  $B$ . Similarly, no vertex of  $B$  is complete to  $A$ .  $\square$

We now argue that we can focus on a specific type of proper  $W$ -joins. A  $W$ -join  $(A, B)$  is *partitionable* if there are partitions of  $A$  into non-empty sets  $A', A''$  and of  $B$  into non-empty sets  $B', B''$  such that  $A'$  is anticomplete to  $B''$  and  $B'$  is anticomplete to  $A''$ . A proper  $W$ -join  $(A, B)$  is *shatterable* if it is partitionable with sets  $A', A'', B', B''$  and one of  $(A', B'), (A'', B'')$  is also a proper  $W$ -join; we say it is *unshatterable* otherwise.

**Lemma 13.** *Let  $G$  be a graph with distinct neighbourhoods and let  $(A, B)$  be a proper  $W$ -join in  $G$ . If  $(A, B)$  is partitionable and unshatterable, then  $G[A \cup B]$  is isomorphic to  $C_4$ .*

**Proof.** Suppose  $(A, B)$  is partitionable with sets  $A', A'', B', B''$ . By definition, we have  $|A'|, |A''), |B'|, |B''| \geq 1$ . Without loss of generality,  $|A'| + |B'| \geq |A''| + |B''|$ . Since  $(A, B)$  is unshatterable,  $(A', B')$  is not a proper  $W$ -join. Suppose  $|A'| + |B'| > 2$ . Then, as  $G$  has distinct neighbourhoods,  $(A', B')$  is a  $W$ -join. Consequently,  $(A', B')$  is a proper  $W$ -join by Lemma 12, which is a contradiction. Hence,  $|A'| = |B'| = 1$ . It follows that  $|A''| = |B''| = 1$ . Since  $(A, B)$  is proper,  $G[A \cup B]$  is isomorphic to  $C_4$ .  $\square$

**Lemma 14.** *Let  $G$  be a claw-free graph that is not cobipartite, has distinct neighbourhoods, and has diameter 2. Let  $(A, B)$  be a proper  $W$ -join in  $G$  that is unshatterable. If  $G$  admits a disconnected cut, then there exists a disconnected partition  $V_1, V_2, V_3, V_4$  of  $G$  such that  $V_i \cap (A \cup B) = \emptyset$  for some  $i \in \{1, 2, 3, 4\}$ .*

**Proof.** Let  $V_1, V_2, V_3, V_4$  be any disconnected partition of  $G$  and suppose that  $V_i \cap (A \cup B) \neq \emptyset$  for each  $i \in \{1, 2, 3, 4\}$ . Note that  $V_1$  is anticomplete to  $V_3$  and  $V_2$  is anticomplete to  $V_4$  by definition. Then we may assume without loss of generality that  $V_1 \cap A \neq \emptyset$  and  $V_2 \cap A \neq \emptyset$ . The former implies that  $V_3 \cap (A \cup B) \subseteq B$  and thus  $V_1 \cap (A \cup B) \subseteq A$ . The latter implies that  $V_4 \cap (A \cup B) \subseteq B$  and thus  $V_2 \cap (A \cup B) \subseteq A$ . It follows that  $(A, B)$  is partitionable with sets  $V_1 \cap A, V_2 \cap A, V_3 \cap B, V_4 \cap B$ . Then, by Lemma 13 and the assumption that  $(A, B)$  is unshatterable,  $G[A \cup B]$  is isomorphic to a  $C_4$ . For  $i \in \{1, 2, 3, 4\}$ , let  $v_i$  be the single vertex in  $V_i \cap (A \cup B)$ , so, for  $i \in \{1, 2, 3, 4\}$ ,  $v_i$  is adjacent to  $v_{i+1}$  with  $v_5 = v_1$ , and moreover,  $A = \{v_1, v_2\}$  and  $B = \{v_3, v_4\}$ .

Let  $P = N(A) \setminus N[B]$ ,  $Q = N(B) \setminus N[A]$ ,  $M = N[A \cup B] \setminus (P \cup Q)$ , and  $R = V(G) \setminus (P \cup Q \cup M)$ . Note that  $P$  is complete to  $A$  and anticomplete to  $B$ , whereas  $Q$  is complete to  $B$  and anticomplete to  $A$ . Moreover,  $M$  is complete to  $A \cup B$ , whereas  $R$  is anticomplete to  $A \cup B$ . This means that  $P \subseteq V_1 \cup V_2$ , because its neighbourhood includes  $v_1$  and  $v_2$ . Moreover,  $Q \subseteq V_3 \cup V_4$ , because its neighbourhood includes  $v_3$  and  $v_4$ , whereas  $M = \emptyset$ , because its neighbourhood includes  $v_1, v_2, v_3$ , and  $v_4$ . Moreover,  $G[P]$  is a clique, because two non-adjacent vertices  $u, v \in P$  with  $v_1$  and  $v_4$  yield a claw. Similarly,  $G[Q]$  is a clique.

Observe that one of the following moves yields the requested disconnected partition of  $G$ :

- moving  $v_1$  to  $V_2$  and  $v_4$  to  $V_3$ , unless  $|V_1| = 1$  or  $|V_4| = 1$ ;
- moving  $v_2$  to  $V_1$  and  $v_3$  to  $V_4$ , unless  $|V_2| = 1$  or  $|V_3| = 1$ .

The crux is to show that  $|V_1|, |V_4| > 1$  or  $|V_2|, |V_3| > 1$ . We now go through several cases.

Suppose that  $R = \emptyset$ . Since  $M = \emptyset$ , it follows that  $P \cup A$  and  $Q \cup B$  forms a cobipartition of  $G$ , a contradiction. Hence,  $R \neq \emptyset$ .

Suppose that  $P = \emptyset$ . Since  $R \neq \emptyset$ , let  $v \in R$ . Note that every path from  $v$  to  $v_1$  must intersect both  $Q$  and  $B$ , two disjoint sets. This contradicts the assumption that  $G$  has diameter 2. Hence,  $P \neq \emptyset$ . Similarly, we derive that  $Q \neq \emptyset$ .

Without loss of generality, assume that  $P \cap V_1 \neq \emptyset$  and let  $x \in P \cap V_1$ . Suppose that  $|V_4| = 1$ , and thus  $V_4 = \{v_4\}$ . Then  $Q \subseteq V_3$ . Now note that, because  $R \neq \emptyset$ , there exists a  $j \in \{1, 2, 3\}$  such that  $R \cap V_j \neq \emptyset$ . We consider all three cases.

Suppose that  $R \cap V_1 \neq \emptyset$ . Since  $V_1$  is anticomplete to  $V_3$ , any path from a vertex in  $R \cap V_1$  to  $v_4$  must contain either at least one internal vertex in  $R \cup P$  and one in  $Q$ , or at least one internal vertex in  $P$  in addition to  $v_1$ . This contradicts the assumption that  $G$  has diameter 2.

Suppose that  $R \cap V_2 \neq \emptyset$ . Then  $|V_2| > 1$ . Since  $Q \neq \emptyset$  and  $Q \subseteq V_3$ , it follows that  $|V_3| > 1$ . Then moving  $v_2$  to  $V_1$  and  $v_3$  to  $V_4$  yields a disconnected partition as requested.

Suppose that  $R \cap V_3 \neq \emptyset$ . Let  $v \in R \cap V_3$ . Then a shortest path from  $v$  to  $v_1$ , which has length 2, cannot intersect both  $Q$  and  $B$ . Hence, any such shortest path must intersect  $P$ . Since  $V_3$  is anticomplete to  $V_1$ , it follows that  $V_2 \cap P \neq \emptyset$ . Hence,  $|V_2| > 1$  and  $|V_3| > 1$ , and then moving  $v_2$  to  $V_1$  and  $v_3$  to  $V_4$  yields a disconnected partition as requested.

We may thus assume that  $|V_4| > 1$ . Since  $|V_1| > 1$ , by moving  $v_1$  to  $V_2$  and  $v_4$  to  $V_3$ , we obtain a disconnected partition as requested.  $\square$

Let  $(A, B)$  be a proper  $W$ -join of a graph  $G$ . For any two adjacent vertices  $a \in A$  and  $b \in B$ , let  $G_{ab}$  be the graph obtained from  $G$  by removing  $A \setminus \{a\}$  and  $B \setminus \{b\}$ . Observe that the graph  $G_{ab}$  is the same regardless of the choice of  $a, b$ .

**Lemma 15.** *Let  $G$  be a claw-free graph that is not cobipartite, has distinct neighbourhoods, and has diameter 2. Let  $(A, B)$  be a proper  $W$ -join of  $G$  that is unshatterable. Then  $G$  admits a disconnected cut if and only if  $G_{ab}$  admits a disconnected cut for any two adjacent vertices  $a \in A$  and  $b \in B$ .*

**Proof.** First suppose that  $G_{ab}$  admits a disconnected partition  $V_1, V_2, V_3, V_4$  for any two vertices  $a, b$ . Let  $a \in V_i$  and  $b \in V_j$  for  $i, j \in \{1, 2, 3, 4\}$ . Then the sets  $V'_1, V'_2, V'_3, V'_4$  obtained from  $V_1, V_2, V_3, V_4$  by adding  $A \setminus \{a\}$  to  $V_i$  and  $B \setminus \{b\}$  to  $V_j$  is a disconnected partition of  $G$ .

Now suppose that  $G$  admits a disconnected cut. Let  $V_1, V_2, V_3, V_4$  be a disconnected partition of  $G$ . By Lemma 14, we may assume without loss of generality that  $V_4 \cap (A \cup B) = \emptyset$ . Note that  $A$  is a clique in  $G$  and  $V_1$  is anticomplete to  $V_3$ , and thus  $A \subseteq V_1 \cup V_2$  or  $A \subseteq V_2 \cup V_3$ . We assume the former without loss of generality. Among all such disconnected partitions, we will assume that  $V_1, V_2, V_3, V_4$  was chosen to minimize  $|A \cap V_1|$ .

We consider several cases. In each of these cases we find two vertices  $a, b$  for which we can construct a disconnected partition of  $G_{ab}$ . Note that this suffices to prove the statement, as the graph  $G_{ab}$  is the same regardless of the choice of  $a, b$ .

First assume that  $A \subseteq V_1$ . Since no vertex of  $B$  is anticomplete to  $A$  by the definition of a proper  $W$ -join and  $V_1$  is anticomplete to  $V_3$ , it follows that  $B \subseteq V_1 \cup V_2$ . Now if  $B \subseteq V_1$ , then let  $a \in A$  and  $b \in B$  be arbitrary adjacent vertices (these exist by the definition of a  $W$ -join) and  $V_1 \setminus ((A \setminus \{a\}) \cup (B \setminus \{b\})), V_2, V_3, V_4$  is a disconnected partition of  $G_{ab}$ . Otherwise, let  $b \in B \cap V_2$  and let  $a$  be an arbitrary vertex of  $A$  that is adjacent to  $b$  (which exists by the definition of a proper  $W$ -join). Then  $V_1 \setminus ((A \setminus \{a\}) \cup B), V_2 \setminus (B \setminus \{b\}), V_3, V_4$  is a disconnected partition of  $G_{ab}$ .

Now assume that  $A \subseteq V_2$ . Note that  $B \subseteq V_1 \cup V_2 \cup V_3$ . Since  $B$  is a clique and  $V_1$  is anticomplete to  $V_3$ , it follows that  $B \subseteq V_1 \cup V_2$  or  $B \subseteq V_2 \cup V_3$ . First, assume that  $B \subseteq V_2$ . Let  $a \in A$  and  $b \in B$  be arbitrary adjacent vertices; note that  $a, b \in V_2$ . Then  $V_1, V_2 \setminus ((A \setminus \{a\}) \cup (B \setminus \{b\})), V_3, V_4$  is a disconnected partition of  $G_{ab}$ . So we may assume that  $B \not\subseteq V_2$ . Then  $B \cap V_1 \neq \emptyset$  or  $B \cap V_3 \neq \emptyset$ . Without loss of generality, we assume it is the former. Let  $b \in B \cap V_1$  and let  $a \in A$  be any neighbour of  $b$ . Note that  $a \in V_2$ . Then  $V_1 \setminus (B \setminus \{b\}), V_2 \setminus ((A \setminus \{a\}) \cup B), V_3, V_4$  is a disconnected partition of  $G_{ab}$ .

It remains to consider the case where  $A \cap V_1 \neq \emptyset$  and  $A \cap V_2 \neq \emptyset$ . Let  $P = N(A) \setminus N[B]$ ,  $Q = N(B) \setminus N[A]$ ,  $M = N[A \cup B] \setminus (P \cup Q)$ , and  $R = V(G) \setminus (P \cup Q \cup M)$ . Note that  $P$  is complete to  $A$  and anticomplete to  $B$ , whereas  $Q$  is complete to  $B$  and anticomplete to  $A$ . Moreover,  $M$  is complete to  $A \cup B$ , whereas  $R$  is anticomplete to  $A \cup B$ . Then, by the assumptions of the case, we have that  $P \subseteq V_1 \cup V_2$ . Note that  $B \subseteq V_1 \cup V_2 \cup V_3$ . Since  $B$  is a clique and  $V_1$  is anticomplete to  $V_3$ , it follows that  $B \subseteq V_1 \cup V_2$  or  $B \subseteq V_2 \cup V_3$ . Moreover, as  $A \cap V_1 \neq \emptyset$ , it follows from the definition of a proper  $W$ -join that  $B \not\subseteq V_3$ . We now prove that  $B \subseteq V_1 \cup V_2$ .

For contradiction, assume that  $B \cap V_3 \neq \emptyset$  and thus  $B \cap V_2 \neq \emptyset$ . As  $M$  is complete to  $A$  and  $B$  and  $A \cup B$  has a nonempty intersection with each of  $V_1, V_2, V_3$ , it follows from the definition of a disconnected partition that  $M \subseteq V_2$ . Similarly, we derive that  $Q \subseteq V_2 \cup V_3$ ; recall also that  $P \subseteq V_1 \cup V_2$ . Suppose  $V_1 \setminus A \neq \emptyset$ . Then  $V_1 \setminus A, V_2 \cup A, V_3, V_4$  is also a disconnected partition of  $G$ , contradicting our choice of the disconnected partition  $V_1, V_2, V_3, V_4$ , which minimizes  $|A \cap V_1|$ . Hence,  $V_1 \setminus A = \emptyset$  and thus,  $V_1 \subseteq A$ . Then  $P \subseteq V_2$ . By the definition of a  $W$ -join, any path of length 2 from a vertex in  $R$  to a vertex in  $A$  must intersect  $P$  or  $M$ . As  $M \cup P \subseteq V_2$  and  $V_4$  is anticomplete to  $V_2$ , we obtain  $R \cap V_4 = \emptyset$ . Since  $A \cup B \cup P \cup M \cup Q \cup R = V(G)$  and none of  $A, B, P, M, Q, R$  intersects  $V_4$ , it follows that  $V_4 = \emptyset$ , a contradiction.

We may thus assume that  $B \subseteq V_1 \cup V_2$ . First, assume that there exist adjacent vertices  $a \in A$  and  $b \in B$  such that  $|V_1 \cap \{a, b\}| = 1$  (and thus  $|V_2 \cap \{a, b\}| = 1$ ). Then  $V_1 \setminus ((A \cup B) \setminus \{a, b\}), V_2 \setminus ((A \cup B) \setminus \{a, b\}), V_3, V_4$  is a disconnected partition of  $G_{ab}$ . Hence, we may assume that no such two vertices exist. It follows that neither  $B \subseteq V_1$  nor  $B \subseteq V_2$ ; otherwise, such  $a$  and  $b$  would exist by the definition of a proper  $W$ -join. Then we may conclude that  $(A, B)$  is partitionable with sets  $A \cap V_1, A \cap V_2, B \cap V_1, B \cap V_2$ . Since  $(A, B)$  is unshatterable, it follows from Lemma 13 that  $G[A \cup B]$  is isomorphic to  $C_4$ . Hence,  $|A| = |B| = 2$  and  $V_1, V_2$  each contain exactly one vertex of  $A$  and exactly one vertex of  $B$ . Since  $G$  is not cobipartite and  $G$  is connected (as  $G$  has diameter 2), it follows that one of  $P, M, Q$  is non-empty. However, each vertex in  $P \cup M \cup Q$  is adjacent to a vertex of  $V_1$  and a vertex of  $V_2$ . Hence,  $P \cup M \cup Q \subseteq V_1 \cup V_2$ . Without loss of generality,  $(P \cup M \cup Q) \cap V_1 \neq \emptyset$ . Let  $a$  be the single vertex of  $A \cap V_2$  and let  $b$  be the single vertex of  $B \cap V_2$ . Then  $V_1 \setminus (A \cup B), V_2, V_3, V_4$  is a disconnected partition of  $G_{ab}$ . The lemma follows.  $\square$

In Section 5.2 we will show that by iterating the above lemma, we can remove all  $W$ -joins from an input claw-free graph of diameter 2. However, to this end, it is crucial to have a polynomial-time algorithm that actually finds an unshatterable proper  $W$ -join (if it exists). Our algorithm for this problem relies on the  $O(n^2m)$ -time algorithm by King and Reed [26] to find a proper  $W$ -join  $(A, B)$ . We test in linear time whether the proper  $W$ -join is partitionable by considering the graph  $H$  obtained from  $G[A \cup B]$  by removing all edges with both endpoints in  $A$  or in  $B$ . We argue that we can recurse on a smaller proper  $W$ -join if  $H$  has two or more connected components, and that  $(A, B)$  is unshatterable otherwise.

**Lemma 16.** Let  $G$  be a graph with distinct neighbourhoods. Then in  $O(n^2m)$  time, we can find an unshatterable proper  $W$ -join in  $G$ , or report that  $G$  has no proper  $W$ -join.

**Proof.** King and Reed [26] proved that in  $O(n^2m)$  time, one can find a proper  $W$ -join in  $G$ , or report that  $G$  does not admit a proper  $W$ -join. In the former case, let  $(A, B)$  be the proper  $W$ -join that is found. If  $G[A \cup B]$  is isomorphic to  $C_4$ , which can be checked in constant time, then  $(A, B)$  is unshatterable, and we return  $(A, B)$ . So assume otherwise. We can test in linear time whether  $(A, B)$  is partitionable as follows. Let  $H$  be the graph obtained from  $G[A \cup B]$  by removing all edges with both endpoints in  $A$  or in  $B$ . Note that  $H$  is bipartite. Observe also that  $H$  cannot have any singleton connected components, because such a vertex (say  $a \in A$ ) would be anticomplete to the other side ( $B$  in this case), a contradiction to the definition of a proper  $W$ -join. Now note that  $(A, B)$  is partitionable if and only if  $H$  has two or more connected components. If  $H$  has one connected component, which can be tested in linear time, then  $(A, B)$  is not partitionable and thus unshatterable. If  $H$  has three or more connected components, then let  $C$  be any such connected component. Then  $(A, B)$  is partitionable with  $V(C) \cap A, V(C) \cap B, (V(H) \setminus V(C)) \cap A, (V(H) \setminus V(C)) \cap B$ , and since  $H$  does not have any singleton components,  $|V(H) \setminus V(C)| > 3$ , and  $G$  has distinct neighbourhoods,  $(V(H) \setminus V(C)) \cap A, (V(H) \setminus V(C)) \cap B$  is a  $W$ -join in  $G$ . It follows from Lemma 12 that it must be a proper  $W$ -join. Now repeat the algorithm for  $(V(H) \setminus V(C)) \cap A, (V(H) \setminus V(C)) \cap B$ . If  $H$  has exactly two connected components, then let  $C$  be a connected component of  $H$  with the most vertices. Since  $G[A \cup B]$  is not isomorphic to  $C_4$ , it follows that  $C$  has at least three vertices. Then, using that  $G$  has distinct neighbourhoods,  $(V(C) \cap A, V(C) \cap B)$  is a  $W$ -join in  $G$ , which by Lemma 12 must be a proper  $W$ -join. Now repeat the algorithm for  $(V(C) \cap A, V(C) \cap B)$ . This gives the required algorithm.

Observe that the algorithm can recurse at most  $n$  times, since in each recursion step it considers a strictly smaller proper  $W$ -join. Each recursion step takes linear time by performing a breadth-first search on the graph  $H$ . Hence, the running time of our algorithm is dominated by the initial call to the algorithm, which takes  $O(n^2m)$  time.  $\square$

## 5.2. Structure of claw-free graphs and solving disconnected cut

We now show a decomposition of claw-free graphs of diameter 2. In order to do so we need a number of definitions, which all originate in Chudnovsky and Seymour [6], but are reformulated in the style of Hermelin et al. [20,21]. A *trigraph* is defined by a set of vertices and an adjacency relation where any two vertices are either strongly adjacent, semi-adjacent, or strongly anti-adjacent, and every vertex is semi-adjacent to at most one vertex. One may think of a trigraph as a normal graph where some edges are simultaneously present and non-present. In particular, a trigraph without semi-adjacent pairs of vertices is just a graph. We call vertices  $u, v$  of a trigraph *adjacent* if they are strongly adjacent or semi-adjacent, and *anti-adjacent* if they are strongly anti-adjacent or semi-adjacent. We call two sets  $X, Y$  (*strongly*) *complete* if each pair of vertices  $v \in X, w \in Y$  is (strongly) adjacent, and (*strongly*) *anti-complete* if each pair of vertices  $v \in X, w \in Y$  is (strongly) anti-adjacent.

A graph  $H$  is a *thickening* of a trigraph  $G$  if there is a partition of  $V(H)$  into non-empty sets  $X_v$  for each  $v \in V(G)$  such that:

- $X_v$  is a clique for each  $v \in V(G)$ ;
- if  $v, w$  are strongly adjacent in  $G$ , then  $X_v$  is complete to  $X_w$ ;
- if  $v, w$  are strongly anti-adjacent in  $G$ , then  $X_v$  is anti-complete to  $X_w$ ;
- if  $v, w$  are semi-adjacent in  $G$ , then  $X_v$  is neither complete nor anti-complete to  $X_w$ .


A pair of vertices  $v, w$  in a graph  $H$  form *twins* if  $N[v] = N[w]$ .

Let  $H$  be a thickening of a trigraph  $G$ . If  $v \in V(G)$  is not semi-adjacent to another vertex, then the vertices in  $X_v$  form twins. Furthermore, if  $v, w$  are semi-adjacent, then  $(X_v, X_w)$  is a  $W$ -join in  $G$  (recall that  $v$  and  $w$  are not semi-adjacent to any other vertices in  $G$ ). In particular, if  $H$  contains neither twins nor  $W$ -joins, then  $G$  and  $H$  are isomorphic.

A *strip-structure* of a connected graph  $G$  consists of a connected multigraph  $H$  (with parallel edges and self-loops), a nonempty set  $X_e \subseteq V(G)$  for each  $e \in E(H)$ , and a nonempty set  $X_{e,y} \subseteq X_e$  for each  $e \in E(H)$  and  $y \in V(H)$  such that  $e$  is incident to  $y$ , such that

- the sets  $X_e$  partition  $V(G)$ ;
- for each  $e \in E(H)$  incident with two vertices  $y, y' \in V(H)$ , each vertex in  $X_{e,y} \cap X_{e,y'}$  is anti-complete to  $X_e \setminus (X_{e,y} \cup X_{e,y'})$ ;
- for each  $y \in V(H)$ , the graph induced by the union, over all  $e \in E(H)$  incident to  $y$ , of the sets  $X_{e,y}$  is a clique;
- if  $v, w$  are adjacent in  $G$ , then either  $v, w \in X_e$  for some  $e \in E(H)$  or there exist  $e, e' \in E(H)$  incident with the same vertex  $y \in V(H)$  for which  $v \in X_{e,y}$  and  $w \in X_{e',y}$ .

For each  $e \in E(H)$ , the *strip corresponding to  $e$*  is a pair  $(J, Z)$ , where  $Z$  is a set of new vertices, one for each vertex  $y \in V(H)$  incident with  $e$ , and the graph  $J$  is obtained from  $G[X_e]$  by adding  $Z$  and for each  $z \in Z$ , making  $z$  complete to  $X_{e,y}$ , where  $y \in V(H)$  is the vertex corresponding to  $z$ . The definition of a strip-structure implies that each strip  $(J, Z)$  has  $|Z| = 1$  (if  $e$



**Fig. 2.** This figure is inspired by [20, Fig. 1]. The left figure is a claw-free graph  $G$ . A strip-structure is indicated by the coloured ovals. The middle figure shows the graph  $H$  underlying the strip-structure; the edge  $b$  indicates a self-loop. The lighter-coloured ovals in the left figure show the sets  $X_e$  for  $e \in E(H)$ . The darker-coloured ovals show the sets  $X_{e,y}$  for each  $e \in E(H)$  and each vertex  $y \in V(H)$  to which  $e$  is incident. The right top figure shows an example of the trivial line graph strip with  $|Z| = 2$ . The right bottom figure shows an example of a stripe with  $|Z| = 2$ . The lighter-coloured ovals show  $X_e = V(J) \setminus Z$  and the darker-coloured ovals show  $X_{e,y}$  for each  $y \in V(H)$  to which  $e$  is incident. Note that strips  $a$ ,  $b$ , and  $f$  in the left figure are trivial line graph strips. Strips  $c$ ,  $d$ , and  $g$  are stripes and might look different depending on  $X_e$ ; the stripe in the right panel corresponds to  $f$  (the stripes corresponding to  $c$  and  $d$  are not pictured). Observe that indeed the set  $Z \subseteq V(J)$  is not part of  $G$ .

is a self-loop) or  $|Z| = 2$  (otherwise). We may think of  $Z$  as being ‘representatives’ of the rest of the graph, but note that the vertices of  $Z$  are not part of  $G$ . We refer to Fig. 2 for an example.

A strip  $(J, Z)$  is a *trivial line graph strip* if  $|Z| = 1$  and  $J$  is a 2-vertex path, or if  $|Z| = 2$  and  $J$  is a 3-vertex path. A strip  $(J, Z)$  is a *stripe* if no vertex in  $V(J) \setminus Z$  is adjacent to more than one vertex of  $Z$ . In particular  $(J, Z)$  is a stripe if  $|Z| = 1$ , and moreover, if  $(J, Z)$  is a stripe with  $|Z| = 2$  for  $e = yy'$ , then  $X_{e,y}$  and  $X_{e,y'}$  are disjoint. See also Fig. 2. A *thickening of a stripe*  $(J, Z)$  is defined as usual, except that it must hold that  $|X_z| = 1$  for each  $z \in Z$ .

We now define an XX-trigraph, XX-graphs, and XX-trigraph stripes. The details are actually unimportant, because we will only use the fact that XX-(tri)graphs (stripes) have at most 13 vertices. Let  $G$  be a trigraph on vertices  $v_1, \dots, v_{13}$  such that  $v_i$  is strongly adjacent to  $v_{i+1}$  for  $i = 1, 2, 3, 4, 5$ ;  $v_6$  is strongly adjacent to  $v_1$ ;  $v_i$  is anti-adjacent to  $v_j$  for  $1 \leq i < j \leq 6$  with  $1 < j - i < 5$ ;  $v_7$  is strongly adjacent to  $v_1$  and  $v_2$ ;  $v_8$  is strongly adjacent to  $v_4, v_5$ , and possibly adjacent to  $v_7$ ;  $v_9$  is strongly adjacent to  $v_1, v_2, v_3$ , and  $v_6$ ;  $v_{10}$  is strongly adjacent to  $v_3, v_4, v_5$ , and adjacent to  $v_9$ ;  $v_{11}$  is strongly adjacent to  $v_1, v_3, v_4, v_6, v_9$ , and  $v_{10}$ ;  $v_{12}$  is strongly adjacent to  $v_2, v_3, v_5, v_6, v_9$ , and  $v_{10}$ ;  $v_{13}$  is strongly adjacent to  $v_1, v_2, v_4, v_5, v_7$ , and  $v_8$ . Then we call  $G - X$  for any  $X \subseteq \{v_7, v_{11}, v_{12}, v_{13}\}$  an *XX-trigraph*. We call  $G - X$  an *XX-graph* if it has no semi-adjacent vertices. We call a stripe  $(J, Z)$  of a strip-structure an *XX-(tri)graph stripe* if  $J$  is an XX-(tri)graph with vertex set  $\{v_1, \dots, v_{13}\} \setminus X$  for some  $X \subseteq \{v_7, v_{11}, v_{12}, v_{13}\}$  such that  $v_7$  and  $v_8$  are strongly anti-adjacent and  $Z = \{v_7, v_8\} \setminus X$ . Note that for any (thickening of an) XX-(tri)graph stripe, it holds that  $V(J) \setminus N[Z] \neq \emptyset$  (take  $v_6$  or  $X_{v_6}$ ).

**Theorem 3** ([21, Theorem 6.8, simplified]). *Let  $G$  be a connected claw-free graph with  $\alpha(G) > 3$  such that  $G$  does not admit twins or proper  $W$ -joins. Then*

- $G$  is a thickening of an XX-trigraph,
- $G$  is a proper circular-arc graph, or
- $G$  admits a strip-structure such that for each strip  $(J, Z)$ 
  - $(J, Z)$  is a trivial line graph strip, or
  - $(J, Z)$  is a stripe for which  $J$  is connected and
    - \*  $\alpha(J) \leq 3$ , and  $V(J) \setminus N_J[Z] \neq \emptyset$ ,
    - \*  $|Z| = 1$  and  $J$  is a proper circular-arc graph,
    - \*  $|Z| = 2$  and  $J$  is a proper interval graph, or
    - \*  $(J, Z)$  is a thickening of an XX-trigraph stripe.

Moreover, we can distinguish the cases and find the strip-structure in polynomial time.

**Theorem 4.** *Let  $G$  be a connected claw-free graph with  $\alpha(G) > 3$  such that  $G$  does not admit twins or  $W$ -joins. Then*

- $G$  is an XX-graph,
- $G$  is a proper circular-arc graph, or
- $G$  admits a strip-structure such that for each strip  $(J, Z)$ 
  - $(J, Z)$  is a trivial line graph strip, or
  - $(J, Z)$  is a stripe for which  $J$  is connected and

- \*  $\alpha(J) \leq 3$ , and  $V(J) \setminus N_J[Z] \neq \emptyset$ ,
- \*  $|Z| = 1$  and  $J$  is a proper circular-arc graph,
- \*  $|Z| = 2$  and  $J$  is a proper interval graph, or
- \*  $(J, Z)$  is an XX-graph stripe.

Moreover, we can distinguish the cases and find the strip-structure in polynomial time.

**Proof.** Observe that if  $G$  does not admit twins nor  $W$ -joins, then thickening operations effectively do nothing, as discussed previously. Hence, any thickening of an XX-trigraph (stripe) is in fact an XX-graph (stripe).  $\square$

We are now ready to prove Theorem 5.

**Theorem 5.** Every claw-free graph  $G$  of diameter 2 with distinct neighbourhoods, no  $W$ -joins,  $\alpha(G) > 3$ , and  $|V(G)| > 13$  is a proper circular-arc graph or a line graph.

**Proof.** Let  $G$  be as in the theorem statement, and assume that  $G$  is not a proper circular-arc graph. In order to prove that  $G$  is indeed a line graph, we apply Theorem 4 to the graph  $G$ . By the assumptions on  $G$ , we obtain a strip-structure consisting of strips of several possible types. We will argue that this strip-structure can be modified so that it contains only trivial line graph strips. This implies that  $G$  is a line graph.

We first claim that if the strip-structure contains a stripe  $(J, Z)$  such that  $V(J) \setminus N_J[Z] \neq \emptyset$  and  $1 \leq |Z| \leq 2$ , then either  $|Z| = 1$  and  $G$  is isomorphic to  $J$  or to  $J - Z$ , or  $|Z| = 2$  and  $G$  is isomorphic to  $J - Z'$  for some  $Z' \subseteq Z$  or to  $J$  where the vertices in  $Z$  have been identified or made adjacent.

Let  $(J, Z)$  be a stripe such that  $V(J) \setminus N_J[Z] \neq \emptyset$  and  $1 \leq |Z| \leq 2$ . Let  $x \in V(J) \setminus N_J[Z]$ . Since  $G$  has diameter 2, every vertex of  $V(G) \setminus (V(J) \setminus Z)$  must be in  $N_G(N_J(z))$ . Now observe that  $N_G(N_J(z)) \setminus V(J)$  is a clique for each  $z \in Z$  by the definition of a strip-structure.

Consider the case  $|Z| = 1$ . Let  $X = N_G(N_J(Z)) \setminus V(J)$ . Note that the neighbourhood of every vertex in  $X$  is  $N_J(Z)$ . Since  $X$  is a clique and  $G$  has distinct neighbourhoods, it follows that  $|X| \leq 1$ . The claim follows.

Consider the case  $|Z| = 2$  and let  $Z = \{z_1, z_2\}$ . Let  $X = N_G(N_J(z_1)) \setminus V(J)$  and  $Y = N_G(N_J(z_2)) \setminus V(J)$ . Suppose that  $|X \cup Y| = |X \cap Y|$ . Then all vertices in  $X \cup Y$  have the same neighbourhood, namely  $X \cup Y \cup N_J(Z)$ , which by the distinct neighbourhood assumption implies that  $|X \cup Y| = 1$ . This implies the claim. A similar line of reasoning applies if  $|X \cap Y| = 0$  and  $X$  and  $Y$  are anticomplete. Otherwise, suppose that  $|X \cup Y| > 2$ . Then let  $A = X \setminus Y$ ,  $B = Y \setminus X$ , and iteratively assign the vertices of  $X \cup Y$  to the smallest of  $A$  and  $B$ . Then  $(A, B)$  is a  $W$ -join in  $G$ , a contradiction. Hence, we may assume that  $|X \cup Y| \leq 2$ . Suppose that  $|X \cup Y| = |X \cap Y| = 2$ . Then both vertices in  $X \cup Y$  have the same neighbourhood, namely  $X \cup Y \cup N_J(Z)$ , a contradiction. Then  $|X \cap Y| = 1$ , which implies the claim, or otherwise,  $|X \cup Y| \leq 2$ , which then also implies the claim.

Now consider the possible stripes in the strip-structure. If  $1 \leq |Z| \leq 2$ ,  $\alpha(J) \leq 3$ , and  $V(J) \setminus N_J[Z] \neq \emptyset$ , then applying the claim, it follows that  $\alpha(G) \leq 3$ , a contradiction. If  $(J, Z)$  is an XX-graph stripe, then  $V(J) \setminus N_J[Z] \neq \emptyset$ , and thus applying the claim,  $G$  has at most 13 vertices, a contradiction.

Suppose that  $|Z| = 1$  and  $J$  is a proper circular-arc graph. If  $V(J) \setminus N_J[Z] = \emptyset$ , then the stripe can be decomposed into  $|V(J)| - 1$  trivial line graph strips, one for each vertex in  $V(J) \setminus Z$ . Otherwise, that is, if  $V(J) \setminus N_J[Z] \neq \emptyset$ , then applying the claim, it follows that  $G$  is a proper circular-arc graph, a contradiction.

Suppose that  $|Z| = 2$  and  $J$  is a proper interval graph. Let  $Z = \{z_1, z_2\}$ . If  $V(J) \setminus N_J[Z] = \emptyset$ , then  $(N_J(z_1), N_J(z_2))$  would induce a  $W$ -join in  $G$ , a contradiction unless  $|N_J(z_1)| = |N_J(z_2)| = 1$ . That implies  $|V(J)| = 4$  and  $J$  is in fact a four-vertex path. Then the stripe can be decomposed into two trivial line graph strips by adding a new node to the strip-structure that supports them both. Otherwise, that is if  $V(J) \setminus N_J[Z] \neq \emptyset$ , then applying the claim,  $G$  is a proper interval graph or a proper circular-arc graph. The latter can be seen from the fact that there trivially exists a representation of  $J$  as a proper interval graph in which the intervals corresponding to  $Z$  extend farthest left and right in the representation. By bending the representation around the circle, we obtain a proper circular-arc graph. In either case, we obtain a contradiction.

From this, it follows that each strip in the (modified) strip-structure must be a trivial line graph strip. This implies that  $G$  is a line graph, as claimed.  $\square$

We now apply Theorem 5 and results from previous sections to obtain our main result.

**Theorem 6.** DISCONNECTED CUT is  $O(n^3m)$ -time solvable for claw-free graphs.

**Proof.** Let  $G$  be a connected claw-free graph on  $n$  vertices and  $m$  edges. We will either find a disconnected cut or conclude that  $G$  has no disconnected cut. Assume  $n \geq 14$ . We compute the diameter of  $G$  in  $O(n^2)$  time. By Lemma 1,  $G$  has no disconnected cut if its diameter is 1 and has a disconnected cut if its diameter is at least 3. Assume the diameter of  $G$  is 2. We check if  $\alpha(G) \leq 3$  in  $O(n(m + n \log n))$  time [12]. If so, then we decide if  $G$  has a disconnected cut in  $O(n^3)$  time by Lemma 7. Assume  $\alpha(G) > 3$ . Hence,  $G$  is not cobipartite.

Next, we check whether  $G$  contains a vertex  $u$  for which there exists a vertex  $v$  such that  $N(u) \setminus \{v\} \subseteq N(v) \setminus \{u\}$ . This takes  $O(n^3)$  time. If so, then we remove  $u$  from  $G$  (and restart the algorithm with the resulting graph, which is still connected and claw-free). This is correct by Lemma 5. Hence, we may assume that  $G$  has distinct neighbourhoods.

Then, we get rid of all  $W$ -joins in  $G$ . Since  $G$  has distinct neighbourhoods, it follows from Lemma 12 that every  $W$ -join in  $G$  is a proper  $W$ -join. Using Lemma 16, in  $O(n^2m)$  time, we can find an unshatterable  $W$ -join in  $G$  or correctly decide that  $G$  does not admit a proper  $W$ -join (and thus no  $W$ -join). In the former case, we apply Lemma 15 to the unshatterable proper  $W$ -join  $(A, B)$  that is found. This takes linear time. We then restart the algorithm on the graph  $G_{ab}$  found by Lemma 15 (note that  $G_{ab}$  is still connected and claw-free). Since  $|A| + |B| \geq 3$ ,  $|V(G_{ab})| < |V(G)|$  and thus we can recurse at most  $n$  times. Hence, we may assume that  $G$  admits no  $W$ -joins.

Next, we check if  $G$  is a circular-arc graph in linear time by Lemma 8. If so, then we apply Theorem 1 to decide if  $G$  has a disconnected cut in  $O(n^2)$  time. Hence, we may assume that  $G$  is not (proper) circular-arc. By Theorem 5 this means that  $G$  is a line graph. Hence, we apply Theorem 2 to decide whether  $G$  admits a disconnected cut in  $O(n^4)$  time. This finishes the description of the algorithm. The running time is clearly  $O(n^3m)$ .  $\square$

Recall from [23] and [16] that  $C_4$ -CONTRACTIBILITY and  $C_4$ -COMPACTION, respectively, are equivalent to DISCONNECTED CUT on graphs of diameter 2. Combining these claims with Theorem 5 leads to the following two consequences.

**Corollary 1.**  $C_4$ -CONTRACTIBILITY is  $O(n^3m)$ -time solvable for claw-free graphs of diameter 2.

**Corollary 2.**  $C_4$ -COMPACTION is  $O(n^3m)$ -time solvable for claw-free graphs of diameter 2.

### 6. H-free graphs for graphs H with at most four vertices

In this section we show that the  $K_4$  is the only 4-vertex graph  $H$  for which the computational complexity of DISCONNECTED CUT is still open on  $H$ -free graphs. To prove this claim, we need to show three additional results.

A graph  $G = (V, E)$  is *complete  $k$ -partite* if  $V$  can be partitioned into  $k$  independent sets  $A_1, \dots, A_k$  for some integer  $k \geq 2$ , such that two vertices are adjacent if and only if they belong to two different sets  $A_i$  and  $A_j$ . The graph  $\overline{P_1 + P_3}$ , which is a triangle with a pendant vertex, is also known as the *paw*. Olariu proved the following result for paw-free graphs.

**Lemma 17 ([31]).** Every connected  $\overline{P_1 + P_3}$ -free graph is either  $C_3$ -free or complete  $k$ -partite for some  $k \geq 3$ .

We also need the following lemma from [16] for  $C_3$ -free graphs (or we could use the more general result that DISCONNECTED CUT is polynomial-time solvable on  $(C_3 + P_1)$ -free graphs in [10]).

**Lemma 18 ([16]).** DISCONNECTED CUT is  $O(n^3)$ -time solvable for  $C_3$ -free graphs.

**Lemma 19.** DISCONNECTED CUT is  $O(n^3)$ -time solvable for  $\overline{P_1 + P_3}$ -free graphs.

**Proof.** Let  $G$  be a  $\overline{P_1 + P_3}$ -free graph. By using brute force we check in  $O(n^3)$  time if  $G$  contains a  $C_3$ . If  $G$  is  $C_3$ -free, then we apply Lemma 18. Otherwise, by Lemma 17,  $G$  is complete  $k$ -partite for some integer  $k \geq 3$ . We claim that  $G$  has no disconnected cut. For contradiction, let  $V_1, V_2, V_3, V_4$  be a disconnected partition of  $G$ . Let  $A_1, \dots, A_k$  be the partition classes of  $G$ . We may assume without loss of generality that  $V_1 \cap A_1 \neq \emptyset$ . Since  $A_1$  is complete to  $A_i$  for each  $i \neq 1$  and  $V_1$  is anticomplete to  $V_3$ , we know that  $V_3 \cap A_i = \emptyset$  for all  $i \neq 1$ . It follows that  $\emptyset \subset V_3 \subseteq A_1$ . By the same argument,  $\emptyset \subset V_1 \subseteq A_1$ . The same line of reasoning implies that  $\emptyset \subset V_i \subseteq A_2$  and  $\emptyset \subset V_i \subseteq A_4$  for some  $i$ , say  $i \in \{1, 2\}$ . Then  $A_3 \cap (V_1 \cup V_2 \cup V_3 \cup V_4) = \emptyset$ , a contradiction.  $\square$

We recall that a biclique is a complete bipartite graph  $K_{r,s}$  for some integers  $r, s \geq 1$ . Recall also that the  $2K_2$ -PARTITION problem is to decide if a graph  $G$  has a  $2K_2$ -partition, or equivalently, if the vertex set of  $G$  can be partitioned into two non-empty sets  $S$  and  $T$  such that  $G[S]$  and  $G[T]$  are bicliques. Moreover, we recall that  $G$  has a  $2K_2$ -partition if and only if  $\overline{G}$  has a disconnected cut. We need the following lemma, which follows from Lemma 7.

**Lemma 20 ([10]).**  $2K_2$ -PARTITION is  $O(n^3)$ -time solvable for  $K_4$ -free graphs.

We can now prove the following lemma.

**Lemma 21.** DISCONNECTED CUT is  $O(n^3)$ -time solvable for  $(P_1 + P_3)$ -free graphs.

**Proof.** Let  $G = (V, E)$  be a  $(P_1 + P_3)$ -free graph on  $n$  vertices. We may assume that  $n \geq 4$ . We check if the  $\overline{P_1 + P_3}$ -free graph  $\overline{G}$  has a  $2K_2$ -partition by applying the following algorithm, which has running time  $O(n^3)$ . If  $\overline{G}$  has more than two

**Table 1**

The complexity of DISCONNECTED CUT on  $H$ -free graphs when  $H$  has four vertices.

$H$	Complexity of DISCONNECTED CUT
$4P_1$	polynomial [10]
$2P_1 + P_2$ (co-diamond)	polynomial [10]
$2P_2$	polynomial [7]
$P_1 + P_3$ (co-paw)	polynomial ( <b>this work</b> : Lemma 21)
$C_3 + P_1$ (co-claw)	polynomial [10]
$K_{1,3}$ (claw)	polynomial ( <b>this work</b> : Theorem 5)
$P_4$	polynomial [16]
$\overline{P_1 + P_3}$ (paw)	polynomial ( <b>this work</b> : Lemma 19)
$2P_1 + P_2$ (diamond)	polynomial ( <b>this work</b> : Lemma 22)
$K_4$	open

disconnected components,  $\overline{G}$  has no  $2K_2$ -partition. Suppose  $\overline{G}$  has exactly two components  $D_1$  and  $D_2$ . Then each  $D_i$  must have at least two vertices and must contain a complete bipartite spanning subgraph. The latter condition is true if and only if each  $\overline{D_i}$  is disconnected. Suppose  $\overline{G}$  has exactly one component. If  $\overline{G}$  is  $C_3$ -free, then we apply Lemma 20. Otherwise,  $\overline{G}$  is complete  $k$ -partite for some  $k \geq 3$  due to Lemma 17. Then, as  $n \geq 4$ , we observe that  $\overline{G}$  has a  $2K_2$ -partition.  $\square$

The graph  $\overline{2P_1 + P_2}$  is also known as the *diamond*.

**Lemma 22.** DISCONNECTED CUT is  $O(n^3)$ -time solvable for  $\overline{2P_1 + P_2}$ -free graphs.

**Proof.** Let  $G$  be a  $\overline{2P_1 + P_2}$ -free graph. We check in  $O(n^2)$  time if  $G$  has a dominating vertex. If so, then  $G$  has no disconnected cut due to Lemma 2. Assume  $G$  has no dominating vertex. As  $G$  is  $\overline{2P_1 + P_2}$ -free, the neighbourhood  $N(u)$  of each vertex must be  $P_3$ -free, and thus  $G[N(u)]$  is the disjoint union of one or more complete graphs. If  $G[N(u)]$  is the disjoint union of at least two complete graphs, then  $u$  has a disconnected neighbourhood. Consequently,  $G$  has a disconnected cut due to Lemma 3. We check in  $O(n^3)$  time if  $G$  has a vertex whose neighbourhood is a disjoint union of at least two complete graphs. Suppose  $G$  has no such vertex, so every neighbourhood  $N(u)$  is a clique. Then  $G$  itself must be a clique and thus  $G$  has no disconnected cut.  $\square$

We are now ready to prove the following summary for DISCONNECTED CUT restricted to  $H$ -free graphs; see also Table 1.

**Theorem 7.** Let  $H \neq K_4$  be a graph on at most four vertices. Then DISCONNECTED CUT is polynomial-time solvable for  $H$ -free graphs.

**Proof.** We may assume that  $H$  has exactly four vertices. Let  $H$  have  $p$  connected components. If  $p = 4$ , then  $H = 4P_1$  and we can use Lemma 7, proven in [10]. If  $p = 3$ , then  $H = 2P_1 + P_2$ . This case was also proven in [10]. If  $p = 2$ , then  $H = 2P_2$ , or  $H = P_1 + P_3$ , or  $H = C_3 + P_1$ . The first case was proven in [7]. The second case is proved in Lemma 21. The third case was proven in [10]. If  $p = 1$ , then  $H = K_{1,3}$ , or  $H = P_4$ , or  $H = \overline{P_1 + P_3}$ , or  $H = \overline{2P_1 + P_2}$ . The first case follows from Theorem 5. The second case follows from a result of [16], which states that DISCONNECTED CUT is polynomial-time solvable for graphs  $G = (V, E)$  with a dominating edge, that is, an edge  $e = xy$  with  $N_G(x) \cup N_G(y) = V$ . Every  $P_4$ -free graph is a cograph (and vice versa). It follows from the definition of a connected cograph that every connected  $P_4$ -free graph on at least two vertices has a spanning complete bipartite subgraph, and thus a dominating edge (take an edge with endpoints in each of the two partition classes of the spanning complete bipartite subgraph). The third case is prove in Lemma 19. The fourth case is prove in Lemma 22.  $\square$

## 7. Open problems

In light of Corollaries 1 and 2 we ask about the complexities of  $C_4$ -CONTRACTIBILITY and  $C_4$ -COMPACTION for claw-free graphs of diameter at least 3. We note that the NP-complete problem  $P_4$ -CONTRACTIBILITY [4] is polynomial-time solvable for claw-free graphs [15].

The complexity classifications of  $H$ -COMPACTION and SURJECTIVE  $H$ -COLOURING are both wide open. In particular, it is not known if there exists a graph  $H$  for which these two problems have a different complexity. However, if we impose restrictions on the input graph, such a graph  $H$  is known:  $C_4$ -COMPACTION is NP-complete for graphs of diameter 3 [33], whereas SURJECTIVE  $C_4$ -COLOURING (being equivalent to DISCONNECTED CUT) is trivial on this graph class. In contrast to claw-free graphs, graphs of diameter 3 do not form a hereditary graph class, that is, they are not closed under vertex deletion. This leads to the natural question if there exist a hereditary graph class  $\mathcal{G}$  and a graph  $H$ , such that  $H$ -COMPACTION and SURJECTIVE  $H$ -COLOURING have different complexity when restricted to  $\mathcal{G}$ . Should  $C_4$ -COMPACTION turn out to be NP-complete for claw-free graphs, then due Theorem 5 and the equivalency between DISCONNECTED CUT and SURJECTIVE  $H$ -COLOURING we can take the class of claw-free graphs as  $\mathcal{G}$  and the graph  $C_4$  as  $H$  to find such a pair  $(\mathcal{G}, H)$ .

As our final open problem, we ask what the complexity of DISCONNECTED CUT is for  $K_4$ -free graphs; recall from Theorem 7 that the  $K_4$  is the only 4-vertex graph  $H$  for which the complexity of DISCONNECTED CUT on  $H$ -free graphs is still open.

### Declaration of competing interest

The authors declare that they have no known competing financial interests or personal relationships that could have appeared to influence the work reported in this paper.

### References

- [1] M.J. Atallah, D.Z. Chen, D.T. Lee, An optimal algorithm for shortest paths on weighted interval and circular-arc graphs, with applications, *Algorithmica* 14 (5) (1995) 429–441.
- [2] M. Bodirsky, J. Kára, B. Martin, The complexity of surjective homomorphism problems – a survey, *Discrete Appl. Math.* 160 (12) (2012) 1680–1690.
- [3] F. Bonomo, G. Oriolo, C. Snels, Minimum weighted clique cover on strip-composed perfect graphs, in: *Proc. WG 2012*, in: *Lecture Notes in Computer Science*, vol. 7551, 2012, pp. 22–33.
- [4] A.E. Brouwer, H.J. Veldman, Contractibility and NP-completeness, *J. Graph Theory* 11 (1) (1987) 71–79.
- [5] D.Z. Chen, D.T. Lee, R. Sridhar, C.N. Sekharan, Solving the all-pair shortest path query problem on interval and circular-arc graphs, *Networks* 31 (4) (1998) 249–258.
- [6] M. Chudnovsky, P.D. Seymour, *The Structure of Claw-Free Graphs*, London Mathematical Society Lecture Note Series, vol. 327, Cambridge University Press, 2005, pp. 153–171.
- [7] K. Cook, S. Dantas, E.M. Eschen, L. Faria, C.M.H. de Figueiredo, S. Klein,  $2K_2$  vertex-set partition into nonempty parts, *Discrete Math.* 310 (6–7) (2010) 1259–1264.
- [8] M. Cygan, M. Pilipczuk, M. Pilipczuk, J.O. Wojtaszczyk, A polynomial algorithm for 3-compatible coloring and the stubborn list partition problem (the stubborn problem is stubborn no more), *SIAM J. Comput.* 41 (4) (2012) 815–828.
- [9] S. Dantas, C.M.H. de Figueiredo, S. Gravier, S. Klein, Finding  $H$ -partitions efficiently, *RAIRO Theor. Inform. Appl.* 39 (1) (2005) 133–144.
- [10] S. Dantas, F. Maffray, A. Silva,  $2K_2$ -partition of some classes of graphs, *Discrete Appl. Math.* 160 (18) (2012) 2662–2668.
- [11] C.M.H. de Figueiredo, The P versus NP-complete dichotomy of some challenging problems in graph theory, *Discrete Appl. Math.* 160 (18) (2012) 2681–2693.
- [12] Y. Faenza, G. Oriolo, G. Stauffer, Solving the weighted stable set problem in claw-free graphs via decomposition, *J. ACM* 61 (4) (2014) 20:1–20:41.
- [13] T. Feder, P. Hell, List homomorphisms to reflexive graphs, *J. Comb. Theory, Ser. B* 72 (2) (1998) 236–250.
- [14] T. Feder, P. Hell, S. Klein, R. Motwani, List partitions, *SIAM J. Discrete Math.* 16 (3) (2003) 449–478.
- [15] J. Fiala, M. Kaminski, D. Paulusma, A note on contracting claw-free graphs, *Discret. Math. Theor. Comput. Sci.* 15 (2) (2013) 223–232.
- [16] H. Fleischner, E. Mujuni, D. Paulusma, S. Szeider, Covering graphs with few complete bipartite subgraphs, *Theor. Comput. Sci.* 410 (21–23) (2009) 2045–2053.
- [17] P.A. Golovach, M. Johnson, B. Martin, D. Paulusma, A. Stewart, Surjective  $H$ -colouring: new hardness results, *Computability* 8 (1) (2019) 27–42.
- [18] F. Harary, *Graph Theory*, Addison-Wesley, Reading MA, 1969.
- [19] P. Hell, Graph partitions with prescribed patterns, *Eur. J. Comb.* 35 (2014) 335–353.
- [20] D. Hermelin, M. Mnich, E.J. van Leeuwen, Parameterized complexity of induced graph matching on claw-free graphs, *Algorithmica* 70 (3) (2014) 513–560.
- [21] D. Hermelin, M. Mnich, E.J. van Leeuwen, G.J. Woeginger, Domination when the stars are out, *ACM Trans. Algorithms* 15 (2) (2019) 25:1–25:90.
- [22] T. Ito, M. Kaminski, D. Paulusma, D.M. Thilikos, On disconnected cuts and separators, *Discrete Appl. Math.* 159 (13) (2011) 1345–1351.
- [23] T. Ito, M. Kaminski, D. Paulusma, D.M. Thilikos, Parameterizing cut sets in a graph by the number of their components, *Theor. Comput. Sci.* 412 (45) (2011) 6340–6350.
- [24] M. Kaminski, D. Paulusma, A. Stewart, D.M. Thilikos, Minimal disconnected cuts in planar graphs, *Networks* 68 (4) (2016) 250–259.
- [25] A.D. King, *Claw-free graphs and two conjectures on  $\omega$ ,  $\Delta$ , and  $\chi$* , PhD thesis, University, Montreal, Canada, 2009.
- [26] A.D. King, B.A. Reed, Bounding  $\chi$  in terms of  $\omega$  and  $\delta$  for quasi-line graphs, *J. Graph Theory* 59 (2008) 215–228.
- [27] S. Mandal, A. Pal, M. Pal, An optimal algorithm to find centres and diameter of a circular-arc graph, *Electron. Int. J. Adv. Model. Optim.* 9 (2007) 155–170.
- [28] B. Martin, D. Paulusma, The computational complexity of disconnected cut and  $2K_2$ -partition, *J. Comb. Theory, Ser. B* 111 (2015) 17–37 (conference version Proc. CP 2011, LNCS 6876, 561–575).
- [29] B. Martin, D. Paulusma, E.J. van Leeuwen, Disconnected cuts in claw-free graphs, in: *Proc. ESA 2018*, in: *Leibniz International Proceedings in Informatics*, vol. 112, 2018, pp. 61:1–61:14.
- [30] R.M. McConnell, Linear-time recognition of circular-arc graphs, *Algorithmica* 37 (2) (2003) 93–147.
- [31] S. Olariu, Paw-free graphs, *Inf. Process. Lett.* 28 (1) (1988) 53–54.
- [32] R.B. Teixeira, S. Dantas, C.M.H. de Figueiredo, The external constraint 4 nonempty part sandwich problem, *Discrete Appl. Math.* 159 (7) (2011) 661–673.
- [33] N. Vikas, Computational complexity of compaction to reflexive cycles, *SIAM J. Comput.* 32 (1) (2002) 253–280.
- [34] N. Vikas, Algorithms for partition of some class of graphs under compaction and vertex-compaction, *Algorithmica* 67 (2) (2013) 180–206 (conference version Proc. COCOON 2011, LNCS 6842, 319–330).